

AEWA EUROPEAN GOOSE MANAGEMENT PLATFORM



**10th MEETING OF THE
AEWA EUROPEAN GOOSE MANAGEMENT
INTERNATIONAL WORKING GROUP**
16-18 June 2026, Skövde, Sweden



EGMP POPULATION STATUS AND OFFTAKE ASSESSMENT REPORT 2026

*Prepared by the EGMP Data Centre
with contributions from the Dutch Modelling Consortium*

Executive Summary

This report provides the 2026 status, offtake assessment and management guidance for the goose populations managed under the EGMP. The information covers aspects related to population status, survival, productivity, as well as assessment of cumulative impact of derogation and legal hunting and, for some populations, management recommendations.

Pink-footed Goose – Svalbard population

In 2023, the Data Centre committed to investigate potential biases in the biannual counts by exploring the use of GPS-tagged birds to estimate detection probabilities. Since then, detection probabilities have been estimated for the November 2022-2024 and May 2023-2026 censuses. Combined GPS tallies from the November counts revealed a detection probability of 0.84 (sd = 0.04), indicating a negative bias in autumn counts. For spring counts, the locations of GPS-tagged individuals were provided to the observers in Trøndelag, Norway in May 2024, 2025, and 2026 to increase the probability of a complete census, whereas observers were not provided this information in 2023. The detection probability for May 2023 was 0.82 (sd = 0.08), again indicating a negative bias, whereas the detection probability for May 2024-2026 (combined) was 1.03 (sd = 0.05), indicating a positive bias. The updated IPM for Pink-footed Goose now accounts for these biases by drawing year-specific detection probabilities from these distributions for all November counts and all May counts prior to 2024, whereas the May 2024, 2025, and 2026 counts were corrected for by drawing detection probabilities from their pooled distribution of detection rates. Beginning with the May 2025 estimate of 78,749 (63,873 – 96,605), the population grew to an estimated 103,715 (85,569 – 123,544) birds in November 2025. The estimate of the May 2026 population size is 79,602 (60,374 – 102,021), well above the new target of 72,000. Harvests and harvest rates were increasing prior to the implementation of the adaptive harvest management program in 2013 but have been somewhat stable since. Estimates of annual survival have generally decreased during the entire period of record, although there is quite a bit of uncertainty associated with the estimates in the last few years (due to the cessation of the capture-mark-recapture program). The suggested harvest quota for the 2026/2027 hunting season, based on the estimated population of 79,602 individuals and 19 days above freezing in Svalbard in May 2026 is the realistic maximum of 20,000 individuals. For comparison, the realized harvest averaged 9,903 (sd = 824) during the last three years of available data. The 2026 harvest quotas for Norway and Denmark this year are thus 6,000 and 14,000, respectively.

Taiga Bean Goose - Scandinavia (br) population

In the Scandinavia (br) population, winter counts in Denmark (585 individuals) and Scotland (173 individuals) resulted in a total of 758 geese which is less than in previous years. The relatively large number of Taiga Bean Geese observed outside the traditional range in England are assumed to belong to the Finland & NW Russia (br) population. To avoid jeopardizing this population, it is strongly encouraged to improve monitoring efforts, maintain harvest restrictions and investigate the extent of unintentional and illegal offtake.

Taiga Bean Goose - Finland & NW Russia (br) population

With respect to the Finland and NW Russia (br) population, the integrated population model has generally been successful at mediating all sources of count and harvest data to provide biologically reasonable and robust estimates of population status. However, because of increasing challenges in potential bias in population counts due to potential changes in count methodology and unknown changes in the spatial and temporal distribution of birds, no formal assessment was made in 2025. For 2026, a major potential source of this bias in the distribution of birds in October has been accounted for, but persistent issues may nonetheless remain. Due to low counts in January and March 2026, the population appears to have declined slightly from previous years to an estimate of 75,470 (67,280 – 84,183), which is still above the target of at least 70,000 individuals in spring. It is currently unknown whether these lower January and March counts are due to biological or methodological issues, and management plans should exercise a degree of caution while continued investigation into the causes of these low counts are underway. With this in mind, we propose a continuation of the restricted hunting levels to ensure that the population stays above the target of 70,000 birds.

Taiga Bean Goose - Germany & Poland (nbr) population

Research on this population is ongoing, and some important information updates have been made available for the revised Taiga Bean Goose ISSAP. Population size continues to be estimated at 15,000-20,000 individuals,

although improved monitoring in the Polish wintering areas in recent years indicate that this may be an overestimate. Further improvement of monitoring efforts is encouraged. Harvest restriction should remain in place.

Taiga Bean Goose - Central Asia (nbr) population

This population is small and probably declining, apparently with fragmented range and low survival, and it is proposed to maintain harvest restrictions and enforcement. Improved monitoring in the wintering areas is strongly encouraged.

Greylag Goose – NW Europe (br) population

Despite considerable improvements in data availability, it has still not been possible to establish a dynamic and model-based management at MU level. However, technical progress has been made, including the development of a flyway population model, a utility model used to evaluate various offtake strategies in terms of their ability to meet population targets, and a model for estimating number of breeding pairs from post-breeding counts. In 2022, a post-breeding population of 540,115 individuals was estimated for MU1, resulting in an estimated ~132,000 breeding pairs, and a post-breeding population of 748,110 individuals was estimated for MU2, equivalent to ~180,000 breeding pairs, indicating that both MUs are well above the set targets of 70,000 and 80,000 breeding pairs, respectively. Between 2022 and 2025, population estimates based on post-breeding counts have increased for all Range States conducting such counts. In January 2025, the wintering population was estimated at 802,320 individuals, which also indicates a population size well above the target of approximately 545,000 individuals in winter. With a reported offtake of at least 449,602 Greylag Geese in 2024/2025, we continue to suspect that the reported offtake is biased high. Range States are encouraged to maintain the current harvest levels and improve harvest and derogation data collection schemes while continuing to focus on offtake strategies minimizing the need for spring/summer derogation and keeping in mind the decreasing number of Greylag Geese migrating to the southern part of the range.

Barnacle Goose – Russia (br) population

Assessment of the population status of the Russia/Germany & Netherlands population aims to analyse the cumulative impact of derogation and hunting (where legally allowed) on the status of the flyway population and the status of the three individual management units (MUs) which have been defined, for the period 2005/06-2024/25. Data from field counts as well as posterior estimates from the Integrated Population Model (IPM) indicate a flyway population size of about 1.5 million individuals in January 2025, thus four times the FRP. This estimate is slightly below the estimates from January 2024, and data collected in the last five years suggest that the total flyway population size has not undergone an overall growth recently. The Russian breeding population also experienced very low productivity in 2024. The Russian MU1 has by far the largest share (92%) of the total flyway population size. Converted into breeding pairs, numbers in the Russian MU1 are well beyond (5.5 times) the FRP and also well above the 200% threshold level. The Baltic MU2 in summer 2024 was 1.8 times FRP but has dropped below the threshold of 200% FRP (calling for coordination if significant offtake under derogation affects the local breeding populations in Denmark, Finland or Sweden) as a result of an ongoing decline. In the North Sea MU3, the number of breeding pairs has increased (or recovered) lately and was 1.6x FRP, but numbers are still below the 200% threshold, thus requesting coordination of offtake under derogation between Germany and the Netherlands. Gaps in monitoring effort mainly exist in the summer period, both regarding data on abundance and data on productivity.

Barnacle Goose – E Greenland (br) population

After a peak flyway population of 80,000 in 2006 and 2012, abundance declined to 65,516 (55,558 – 76,459) in March 2026. For much of the period of record, abundance on Islay exceeded that in all other wintering areas, but that pattern has been reversed since 2018. Based on the IPM, the total harvest rate of adults has increased over the period of record, from around 0.01 to a peak of 0.05 (0.04 – 0.07) in 2017. Thereafter, harvest rate declined

to 0.02 (0.01 – 0.03) in 2025. Annual survival rate of adults (including both harvest and natural mortality) declined at the same time harvest rates were increasing, suggesting that harvest may have contributed to the decline in flyway abundance, although poorer than average reproduction could also have played a role. There currently is a 1% probability that the March 2026 population is below the FRP of 54,000. However, because of the proximity of the population to the FRP in recent years, the Adaptive Flyway Management Plan requires continued coordination of offtake between Iceland and Scotland to ensure the population does not fall below the FRP.

Action requested from the EGM IWG:

The EGM IWG is requested to take note of the *EGMP Population Status and Offtake Assessment Report 2026* and provide further guidance to the Secretariat and Data Centre.

Preface

This report provides the 2026 status, offtake assessment and management guidance for the goose populations managed under the EGMP. The information covers aspects related to population status, survival, and productivity, as well as assessment of cumulative impact of derogation and legal hunting and, for some populations, management recommendations.

Compilers

Iben H. Sørensen¹, Ryan R. Germain¹, Kees Koffijberg², Jurrian van Irsel² and Jesper Madsen¹

Recommended citation

Sørensen, I.H., Germain, R.R., Koffijberg, K., van Irsel, J., and Madsen, J. (2026). Population Status and Assessment Report 2026. EGMP Technical Report No. 29. Bonn, Germany

Acknowledgements

Many individuals are involved in the data collection in each Range State. Those listed in the EGMP Database may be the ones delivering data to the EGMP Data Centre, but not necessarily the ones responsible for the actual data collection. We would therefore like to thank the network of national coordinators and all volunteers and agencies who contributed to the population counts, the hunters and wildlife councils who delivered data to different schemes across the ranges of these populations or provided wings of shot birds (see EGMP Database for further details and full [acknowledgements](#)). Furthermore, we also wish to thank the EGMP Task Forces and the EGMP Modelling Consortium for helpful reviews of earlier drafts and the EGMP Range States that contributed to the annual budget of the EGMP Data Centre.

Funding organisations

We are very grateful to:

- All EGMP Range States which have contributed with annual voluntary contributions to the EGMP Data Centre activities.
- Jægernes Naturfond in Denmark for funding development of a flyway-wide decision model for use in investigating population-management strategies for Greylag Geese, as well as the French National Office for Biodiversity and the Dutch Ministry of Agriculture, Nature & Food Quality for funding projects focused on estimates of demographic parameters that will be used in the flyway model and for monitoring of crippling rate.
- A range of national initiatives to improve monitoring programs, particularly the development of monitoring programs to estimate the breeding population of Greylag Geese, in Denmark funded by the Danish Environmental Protection Agency, in Finland funded by the Finnish Wildlife Agency and the Finnish Ministry of Agriculture and Forestry, in Sweden funded by the Swedish Environmental Protection Agency, in Norway funded by the Norwegian Environmental Agency and in France by the National Office for Biodiversity and the Ligue pour la Protection des Oiseaux.
- NatureScot and the Department of Housing, Local Government and Heritage Ireland, for funding development of an integrated population model for the E. Greenland/Scotland & Ireland population of Barnacle Geese and the international census which took place in February 2023.

¹ EGMP Data Centre, Dept. of Ecoscience, Aarhus University, Denmark.

²Sovon, Vogelonderzoek, The Netherlands.

- The Ministry of Agriculture, Nature and Food Quality, the Provinces and BIJ12 in The Netherlands for funding development of an integrated population model for the Russian/Germany and Netherlands population of Barnacle Goose, developing a population model for Greylag Geese, and supporting the collection of monitoring data.
- The Finnish Wildlife Agency and Natural Resources Institute for funding development of an integrated population model for Taiga Bean Goose.
- The Danish Environmental Protection Agency for funding the assessment of potential biases in the population counts of Pink-footed Goose.
- The Norwegian Environment Agency, The County Governor of Vestfold and Telemark and the County Governor of Nordland for contribution to the juvenile assessments of Greylag Geese in Norway.

TABLE OF CONTENTS

PREFACE.....	4
COMPILERS.....	4
RECOMMENDED CITATION.....	4
ACKNOWLEDGEMENTS.....	4
FUNDING ORGANISATIONS.....	4
TABLE OF CONTENTS.....	6
INTRODUCTION	8
1.1 THE ASSESSMENT PROCESSES.....	10
2 MONITORING AND ASSESSMENT METHODS.....	11
2.1 POPULATION SIZE	11
2.2 REPRODUCTION.....	12
2.3 OFFTAKE AND SURVIVAL.....	13
2.4 POPULATION ASSESSMENT METHODS	13
3 RESULTS AND DISCUSSION.....	15
3.1 PINK-FOOTED GOOSE <i>ANSER BRACHYRHYNCHUS</i>	15
3.1.1 RANGE STATES AND MANAGEMENT UNITS	15
3.1.2 POPULATION FRP AND TARGET	16
3.1.3 MANAGEMENT STRATEGIES.....	16
3.1.4 ASSESSMENT PROTOCOL	16
3.1.5 POPULATION STATUS.....	18
3.1.6 MANAGEMENT GUIDANCE.....	22
3.2 TAIGA BEAN GOOSE <i>ANSER FABALIS FABALIS</i>	24
3.2.1 RANGE STATES AND POPULATIONS	24
3.2.2 POPULATION TARGETS.....	24
3.2.3 MANAGEMENT STRATEGIES.....	25
3.2.4 ASSESSMENT PROTOCOL	25
3.2.5 STATUS – SCANDINAVIA (BR) POPULATION	27
3.2.6 STATUS – FINLAND & NORTH-WESTERN RUSSIA (BR) POPULATION.....	29
3.2.7 STATUS – GERMANY & POLAND (NBR) POPULATION.....	33
3.2.8 STATUS – CENTRAL ASIA (NBR) POPULATION	33
3.2.9 MANAGEMENT GUIDANCE.....	34
3.3 GREYLAG GOOSE <i>ANSER ANSER</i>	35
3.3.1 RANGE STATES AND MANAGEMENT UNITS	35
3.3.2 POPULATION FRPS AND TARGETS.....	36
3.3.3 MANAGEMENT STRATEGIES.....	36
3.3.4 ASSESSMENT PROTOCOL	36
3.3.5 POPULATION STATUS.....	38
3.3.6 MANAGEMENT GUIDANCE.....	43
3.4 RUSSIA BREEDING POPULATION OF BARNACLE GOOSE <i>BRANTA LEUCOPSIS</i>	46
3.4.1 RANGE STATES AND MANAGEMENT UNITS	46

3.4.2	POPULATION FRPS AND TARGETS	46
3.4.3	MANAGEMENT STRATEGIES	47
3.4.4	ASSESSMENT PROTOCOL	48
3.4.5	STATUS	49
3.4.6	MANAGEMENT GUIDANCE	56
3.5	EAST GREENLAND BREEDING POPULATION OF BARNACLE GOOSE <i>BRANTA LEUCOPSIS</i>	59
3.5.1	RANGE STATES AND MANAGEMENT UNITS	59
3.5.2	POPULATION FRPS AND TARGETS	59
3.5.3	MANAGEMENT STRATEGIES	59
3.5.4	ASSESSMENT PROTOCOL	60
3.5.5	STATUS	61
3.5.6	MANAGEMENT GUIDANCE	65
	APPENDIX A – DATA OVERVIEW	67
A.1.	PINK-FOOTED GOOSE – SVALBARD (BR) POPULATION	67
A.2.	TAIGA BEAN GOOSE	67
A.3.	GREYLAG GOOSE – NW EUROPE (BR) POPULATION	67
A.4.	BARNACLE GOOSE – RUSSIA BREEDING POPULATION	69
A.5.	BARNACLE GOOSE – EAST GREENLAND BREEDING POPULATION	69
	REFERENCES	71

Introduction

The first international management plan to actively manage a migratory population of waterbirds in Europe was adopted in 2012 and implemented in 2013. The plan was for the Svalbard breeding population of Pink-footed Goose and was based on the concept of adaptive management (AM). AM provides a framework for making objective decisions in the face of uncertainty about an ecological system and the impact of management actions. To reduce this uncertainty and improve management over time, AM relies on an iterative cycle of monitoring, assessment, and decision-making.

In 2013, plans for the first iterative cycle were published in the form of a population status report and a harvest assessment report. In May 2016, the European Goose Management Platform (EGMP) was established, following a resolution adopted by the Meeting of the Parties of the African-Eurasian Migratory Waterbird Agreement (AEWA). The platform functions under the framework of AEWA, which provides for the conservation and sustainable use of the migratory waterbird populations it covers. The platform addresses the conservation and management of declining, as well as growing, goose populations in Europe. This is achieved by a coordinated flyway approach amongst all Range States concerned.

The setup of EGMP benefited from experiences with Svalbard Pink-footed Geese and was initially extended to include Taiga Bean Geese. In 2017, four more populations were added to the EGMP; the NW Europe (br) population of Greylag Goose, as well as the three populations of Barnacle Goose: the Russia (br) population, E Greenland breeding population and the Svalbard breeding population. Following the decision by AEWA MOP8 in 2022 to split Taiga Bean Goose, formerly considered as one population consisting of four management units, into four populations, the EGMP encompasses nine populations. In some populations, management units have been established to delineate subpopulations, which are considered to have their own demography and/or dispersal and thus requiring specific management and conservation approaches. Thus, four goose species and their respective populations and management units are currently part of the EGMP (Table 1-1).

Table 1.1. Overview of populations and Management Units (MUs) covered by the EGMP and relevant documents.

Population	Management/Action Plan (ISSMP/ISSAP)			Adaptive Flyway Management Programme (AFMP) / Implementation Plan		
	Link	Adopted	Expires	Link	Adopted	Expires
Svalbard (br) population of Pink-footed Goose	ISSMP	2025	2037	[DRAFT AFMP]	[2026]	-
Scandinavia (br) population of Taiga Bean Goose	ISSAP	2025	2037	Not developed	-	-
Finland & NW Russia (br) population of Taiga Bean Goose	ISSAP	2025	2037	Not developed	-	-
Germany & Poland (nbr) population of Taiga Bean Goose	ISSAP	2025	2037	Not developed	-	-
C Asia (nbr) population of Taiga Bean Goose	ISSAP	2025	2037	Not developed		
NW Europe (br) population of Greylag Goose consisting of 2 MUs; MU1 (migratory) and MU2 (sedentary)	ISSMP	2018	2031	AFMP	2020	2026
Russia (br) population of Barnacle Goose consisting of 3 MUs; MU1 (Arctic), MU2 (Baltic) and MU3 (North Sea)	ISSMP	2018	2031	AFMP	2020	2026
E Greenland (br) population of Barnacle Goose	ISSMP	2018	2031	AFMP	2020	2026
Svalbard (br) population of Barnacle Goose	ISSMP	2018	2031	Not developed	-	-

This report, together with the [EGMP Database](#), comprises a joint population status and harvest assessment for all populations covered by the EGMP. The EGMP Database provides a shared platform for the most up-to-date monitoring information on each population managed under the EGMP (including data sources), whereas this report focuses on the assessment results and management guidance, to be reviewed at the annual meeting of the International Working Group.

Previous EGMP reports are available at: <https://egmp.aewa.info/resources/publications>.

For populations/species where the cumulative impact of derogation and legal hunting is assessed and/or management guidance provided, input and output files of the assessment runs from previous years are available at: <https://gitlab.com/aewa-egmp>. Most recent files (current assessment) and further details are available from the EGMP Data Centre (egmp@ecos.au.dk).

Information on indicators related to other aspects of the management plans, such as socioeconomic issues and ecosystem services provided by geese, are presented in the Adaptive Flyway Management Programmes (AFMPs) in the annex 'Indicator factsheets'. All AFMPs are available here: <https://egmp.aewa.info/resources/action-and-management-plans-adaptive-flyway-management-programmes>.

1.1 The assessment processes

The assessment process is pictured in Figure 1.1-1 and consist of three steps;

1) *Monitoring.*

Periodic monitoring and other data collection is essential for keeping track of the implementation progress for the EGMP ISSMPs, ISSAPs and AFMPs, not least regarding the process for setting hunting regulations and assessing the impact of derogation. Monitoring data refers to measures of abundance (counts or indices based on samples), data on productivity (counts of young and adults) and survival, and data to describe offtake (either hunting bags or derogation data). Monitoring and data collection are ongoing activities, which take place throughout the year, and are conducted according to agreed protocols. Data from monitoring activities are compiled by the EGMP Data Centre, by Sovon Vogelonderzoek Nederland for the Russia (br) population of Barnacle Goose, and by NatureScot for the E Greenland (br) population of Barnacle Goose. See Appendix A for coverage in each country and population and the [EGMP Database](#) for overview of data.

2) *Assessment.*

The data produced by monitoring provides information to estimate the status of the populations and are used along with other information to evaluate progress towards reaching management objectives, as well as to facilitate learning after decisions are made.

For populations/species where population models have been developed, demographic information like population size, productivity and survival rates are based on model estimates, and updated as new data are received. For populations/species without population models and/or updated data, the most current information received from the range states and their monitoring networks is presented. Due to delays in acquiring certain data, some information presented in this report will differ from that in previous reports and may also be subject to updates in future reports.

For populations/species where only derogation is allowed, the cumulative impact of offtake is assessed through retrospective and prospective analyses, investigating the effect of derogation at the population and at the MU level. The effect of the current level of derogation and environmental variables (e.g., avian influenza) is also projected into the future.

For huntable populations/species, a harvest strategy is derived, and annual management guidance is provided. This happens either through a formal adaptive harvest management process as for Pink-footed Goose, or through consensus on quotas informed by simulations as is done for the Finland & North-western Russia (br) population of Taiga Bean Goose (formerly known as the Central MU).

No reporting is provided for the Svalbard/SW Scotland population of Barnacle Goose.

3) *Decision-making.*

The decision-making process takes place by national representatives at the IWG annual meetings. Decision making at each decision point considers management objectives, resource status, and knowledge about consequences of potential actions. Decisions are then implemented by means of management actions on the ground.




Figure 1.1-1. The EGMP assessment process, including annual activities related to monitoring, assessment and decision-making.

2 Monitoring and Assessment Methods

2.1 Population size

Counts of geese managed under the EGMP are performed at different times throughout the year. The counts can be either total counts or counts collected through a sampling program with the aim of estimating the total population size and/or to monitor a trend.

January census: All goose populations managed under the EGMP are covered by the International Waterbird Census (IWC), which takes place in January and has been implemented in most countries forming part of the respective Eurasian flyways. These counts focus on wetland areas, but in some countries include schemes specifically for geese as well, covering occurrence in farmland areas. Field work is usually carried out by a large network of volunteers during daytime on feeding sites or at dawn/dusk at roost sites, but precise methods, and especially coverage, may vary slightly between countries. In addition, some countries (e.g., The Netherlands, Belgium) account for missing geese in the network of counting sites by estimating missing counts ("imputed") with algorithms that account for the long-term trend and the phenology in similar census areas within the region (Hornman et al. 2021; Onkelinx et al. 2017). That way the data used for trend calculations represent a complete dataset and is not subject to variation in counting effort. Goose counts are collected by national coordinators and reported to Wetlands International who coordinates the IWC (van Roomen et al. 2025).

For several species, the January census provides the best available knowledge on the size of the total flyway population, as it has relatively high coverage in all countries and has been in place since the late 1950s, allowing for analyses of long-term time series (Fox and Leafloor 2018). Also, it takes place towards the end of the hunting season for most species, thus allowing an assessment of the effects of offtake. However, for widely dispersed species like e.g., Greylag Goose, the January census only provides information on the overall trend of the entire flyway population, as coverage is currently regarded too low to assess total population size. Moreover, the January count is not suitable to assess the size and trend for some populations and specific MUs as birds from different MUs mix during winter. For these reasons, specific counts are also organised at other times during the year, in order to assess the size of the respective MU-populations. Under the EGMP, data from the IWC is currently only used directly in the assessment of the NW Europe (br) population of Greylag Goose.

Autumn census: In continental Europe, special population counts have previously been made for all grey geese (*Anser sp.*) in November, as well as in September for Greylag Goose (Madsen et al. 1999). In recent years, most Range States have performed additional counts, in some cases covering the entire wintering season. A general issue with the autumn counts is that for huntable species, the counts are likely to occur after the start of the hunting season, which from a modelling and assessment perspective complicates the assessment process.

Spring census: Counts during spring, just before the assessment process in May/June and after the hunting season, is on the other hand the best time of the year to provide knowledge on the population size of huntable species shortly before breeding. For the Svalbard (br) population of Pink-footed Goose a total count is organized in early May, just before they leave for the breeding areas and are highly concentrated in only a few areas. For the Finland & North-western Russia (br) population of Taiga Bean Goose, a count (in addition to the autumn and mid-winter count) is organized in March, when most of the population is gathered in Sweden and good coverage is possible. To estimate population sizes of breeding waterfowl and wader species, including Greylag Goose, France has recently introduced a spring census which will take place at regular intervals (currently planned for every six years).

Summer census: For populations where management is performed at a MU level (e.g., Greylag Goose and the Russian population of Barnacle Goose), summer is the only period in which the size of the population in each MU can be assessed. Summer counts take place from mid-July to early September, under the assumption that birds from the respective MUs have not yet left the country or can be accounted for. This type of census does not only cover breeding birds and their offspring, but also failed breeders and non-breeders (i.e., all individuals within the respective MU). So, compared to regular breeding bird surveys in spring (delivering number of breeding pairs), they give a more comprehensive account of abundance (expressed in individuals) in the post-breeding period, while the number of breeding pairs must then be calculated from the results of the post-breeding censuses. Summer counts are carried out during daytime and focus on wetlands and waterbodies, which in summer host nearly all birds during daytime. Hence, coverage is regarded as high (usually >90%), but in some large countries (e.g., Norway and Finland) it is a challenge to coordinate such counts, and alternative sampling approaches have been developed (see Sørensen et al. 2024). Data is collected through volunteer networks but with substantial professional input (more so than during winter). In the IPM-framework, for the Russia (br) population of Barnacle Geese, the number of breeding pairs is set as the number of individuals of 2 years and older divided by 2. For the NW Europe (br) population of Greylag Goose, the number of breeding pairs in the spring is calculated separately for the two management units, based on the results of the post-breeding counts, following the methodology described in Sørensen et al. 2024 (further details provided in Johnson et al. 2024).

All data is provided by national coordinators or agencies, but in some specific cases may also rely on published information (see [EGMP Database](#) for details).

2.2 Reproduction

In migratory geese, productivity is typically expressed as the proportion of young in the autumn population and is assessed at the autumn staging and wintering grounds by observing the number of young vs. adults in flocks of geese – also called age-ratio counts. Such age counts have been performed for many European goose populations for several decades by skilled experts, providing a long-term time series of their breeding performance (Madsen, Cracknell, and Fox 1999; Hornman et al. 2024). Counts are usually done in October and November, Greylag Goose is however already assessed during July and August (in some cases in combination with the summer census), as it is otherwise difficult to distinguish juveniles from adults (see Koffijberg 2022). Assessing productivity at the staging and wintering grounds is, however, likely to be affected by several factors as we are compelled to sample from an open population, in which the temporal and spatial age composition can vary, e.g. due to differential migration, mortality and flocking behaviour (Gupte et al. 2019). The effect of such factors has been investigated, with the Svalbard Pink-footed Goose as a case study (Jensen et al. 2023b).

2.3 Offtake and survival

Hunting bags: All range states allowing hunting have harvest monitoring schemes in place; ranging from national harvest data recording across harvest data schemes at regional level(s) to harvest data collection by wildfowling clubs (UK). Data are generally gathered on an annual basis, but often with a time lag in publishing the data. Furthermore, in most countries, data are gathered for each huntable waterbird species. Most countries have legislation that requires harvest bags to be reported by all hunters, with the exception of Sweden, France, UK and Wallonia, Belgium that have no legislation requiring harvest bags to be reported by all hunters. Moreover, in most countries waterbird harvest data are collected for all individual hunters throughout the country, but in some countries, data are only collected for hunting units, or only a sample of hunters is surveyed. Thus, in general there is an absence of harmonisation among the different hunting bag collecting schemes in Europe. Moreover, there is a lack of information on how calculations are made with the local/regional data to produce the national hunting bag statistics. Thus, reliable inference about flyway totals is very difficult to attain (Aubry et al. 2020). Furthermore, it is not always clear whether the national derogation data (see below) are additional to, or included in, the reported hunting data in countries where both hunting and derogation occurs. For some species, bias in hunting bag reporting is suspected (Johnson and Koffijberg 2021). Hunting bag data are available online in the following countries: [Belgium](#), [Denmark](#), [Finland](#), [Germany](#), [Greenland](#), [Iceland](#), [Norway](#) and [Sweden](#) (a link is provided in each country name).

Derogation: EU Member States are obliged to report all derogations to the European Commission in annual derogation reports (according to Article 9 in the Birds Directive (Directive 2009/147/EC of the European Parliament and of the Council of 30 November 2009 on the conservation of wild birds), see EU 2020). However, for a number of Member States, the data is only available after a delay of several years. Furthermore, in some countries this reporting involves several administrative levels and with some uncertainty as to the true number of birds killed. Derogation data are available from the EU Eionet central data repository (https://environment.ec.europa.eu/topics/nature-and-biodiversity/birds-directive_en), but for this report data has also been provided directly by the Range States.

Wings and heads: In Denmark, Iceland, Scotland, and Sweden hunters may, on a voluntary basis, submit wings from shot geese to national wing surveys. These wing samples contribute to the knowledge of the temporal variation in the hunting bag, as well as knowledge of age ratio among shot birds. In Denmark, Sweden, Finland, and Latvia, hunters have also been invited to submit (photos of) heads of shot Bean Geese to the national hunting organisation for sub-species identification to estimate the proportion of Taiga Bean Geese in the hunting bag.

Crippling rate: In several goose species, X-ray images have been used to assess the proportion with embedded shotgun pellets (Noer et al. 2007). The incidence of embedded shotgun pellets is an expression of hunting exposure and also plays an important role in the ISSMP/AFMP process from an ethical viewpoint and as they are sub-lethal injuries potentially affecting fitness of the geese. Crippling rate is defined here as the proportion of individuals with at least one embedded shotgun pellet, assessed by processing of X-ray images. Whereas the crippling ratio is the crippling rate divided by the harvest rate. Harvest rate is defined as the proportion of the population being shot (Clausen et al. 2017). In general, there is a need for standardized crippling assessment, which is in progress among those institutes collecting data.

2.4 Population assessment methods

Integrated population models (IPM) are currently used to derive estimates of abundance and demographic rates for four goose populations covered by the EGMP: Svalbard Pink-Footed Goose (Johnson et al. 2020a), the Finland & North-western Russia (br) population of Taiga Bean Goose (Johnson et al. 2020b), E. Greenland Barnacle Goose (McIntosh et al. 2021), and the Russian-Germany-Netherlands population of Barnacle Goose (Baveco et al. 2021). IPMs represent an advanced approach to modelling, in which all available demographic data are

incorporated into a single analysis (Schaub and Abadi 2011). IPMs have many advantages over traditional modelling approaches, including the proper propagation of demographic uncertainty, better precision of demographic rates and population size, and the ability to handle missing data and to estimate latent (i.e., unobserved) variables. They also have the capacity to guide the development of effective monitoring programs. IPMs can also be used to derive optimal offtake strategies or to project the future consequences of offtake strategies that have been defined a priori. Finally, use of a Bayesian estimation framework for IPMs provides a natural framework for adaptation, in which demographic parameters can be updated over time based on observations from operational monitoring programs.

Estimates of abundance, survival, and productivity from an IPM are based on the joint statistical likelihood of all the data used in the model. This likelihood is combined with any prior information that may be available to provide what are called posterior estimates of demography. Because the entire historical record of data is always used, all posterior estimates may change slightly each year as new data are added to the historical record. Moreover, posterior estimates from the IPM are unlikely to match perfectly those derived from an independent analysis of an individual source of data. For example, estimates of survival from analysis of capture-mark-recapture (CMR) data are likely to be slightly different than posterior estimates of survival derived from the IPM. This is because the CMR analysis only uses CMR data, whereas the IPM uses the CMR data, plus census data and all other sources of demographic data, to estimate survival. Thus, a great benefit of using the IPM is more reliable estimates of abundance and demography, which better reflect all the demographic information available for a population, and which are not so sensitive to any sources of bias (e.g., which may occur in CMR-data due to neckband loss or differential survival between marked and unmarked birds).

3 Results and Discussion

3.1 Pink-footed Goose *Anser brachyrhynchus*

3.1.1 Range States and Management Units

This chapter compiles monitoring data on the population status of the Svalbard population of Pink-footed Goose, as well as providing guidance for the upcoming hunting season (2026/2027). In 2025, the revised ISSMP for the Pink-footed Goose was adopted by the Parties to AEWa (Madsen et al. 2025a) with a 12-year lifespan (2026-2037). The Range States for this population originally included Norway, Denmark, Belgium and the Netherlands, and has now been expanded to include Finland, Sweden and Russia (Figure 3.1-1). The expansion reflects the recent development of a new migration route through Sweden and Finland with breeding grounds in Novaya Zemlya in northern Russia. This new group consists of at least 4,000 individuals and is expected to increase further, partly due to immigration from the traditional flyway, partly due to high survival and good productivity in the new group (Madsen et al. 2023). The new group is not considered a separate management unit (see Madsen et al. 2025a), as there is continued exchange of individuals between the traditional and the new flyway, and the new migratory pathway continues to develop. In 2026, it is anticipated that an Adaptive Flyway Management Plan will be adopted for the Pink-footed Goose.




Figure 3.1-1. Annual distribution and migration route of Svalbard Pink-footed Goose traditional population (blue) as well as the new breeding group (red; punctuated line shows a route used by some geese in spring).

3.1.2 Population FRP and target

In the revised ISSMP, a Favourable Reference Population size (FRP) of 49,000 individuals in spring was set for the Svalbard population of Pink-footed Goose (Madsen et al. 2025a). Range States agreed on a new population target of 72,000 individuals in spring, thus acknowledging recent changes in the breeding distribution and migration routes of the population while still aiming to reduce agricultural conflicts, particularly in Norway, as well as tundra degradation due to grazing on breeding grounds in Svalbard.

3.1.3 Management strategies

Legal hunting of Svalbard Pink-footed Geese occurs only in Norway and Denmark. A harvest strategy, which is updated each year, prescribes the harvest quota necessary to maintain the population near its target of 72,000 birds. The harvest quota is allocated between Norway (30%) and Denmark (70%) based on historical proportions of the harvest.

3.1.4 Assessment protocol

We used the integrated population model described by Johnson et al. (2020a). Annual changes in population size in May are described by a difference equation:

$$N_{t+1}^M = N_t^M [s_t + r_t \theta_t (1 - v h_t^n - v h_t^d)]$$

where N_t^M is May population size in year t , s_t is the annual survival rate, r_t is the ratio of young of the year to older birds at the start of the hunting season, θ_t is survival from natural causes, h_t^n and h_t^d are per capita harvest rates of birds aged >1 year in Norway and Denmark, respectively, and v is the differential vulnerability of young relative to older birds in the harvest.

Population size in November is a function of population size in May, six months of natural mortality, and the portion of harvest in Denmark occurring prior to November:

$$N_t^N = N_t^M \theta_t^{6/12} [(1 - h_t^n - h_t^d) + r_t (1 - v h_t^n - v h_t^d)]$$

where N_t^N is November population size and h_t^d is the harvest rate of older birds in Denmark prior to November.

Within the IPM, we specified a generalized linear model for reproductive rate (r) using the number of thaw days (D) in May in Svalbard as a covariate: $r_t = \frac{\gamma_t}{(1-\gamma_t)}$, where γ_t is the binomial probability of young, and:

$$\log\left(\frac{\gamma_t}{(1-\gamma_t)}\right) = \beta_0 + \beta_1 D_t$$

Raw data and the results of the 2026 update of the IPM are available from the [EGMP Data Centre](#). We note that hunting bag data for Denmark in 2025 were not available at the time of analyses, and so were drawn from a distribution of potential bag numbers based on the average and standard deviation of the past 4 years, since harvest numbers have declined in recent years and have been relatively stable in Denmark during this period.

Posterior estimates of natural mortality, differential vulnerability of young to harvest, and the regression coefficients expressing the relationship between thaw days and reproductive success were used to derive an optimal harvest policy. We used a computation algorithm known as stochastic dynamic programming (SDP), which can explicitly account for various sources of uncertainty in modelled systems (Marescot et al. 2013).

For computational purposes, the optimal value (V^*) of a management strategy (A) at time t is the maximum (max) of the expectation (E) of the temporal sum of discounted population utilities:

$$V^*(A_t|x_t) = \max_{(A_t|x_t)} E \left[\sum_{\tau=t}^{\infty} \lambda^{\tau} u(a_{\tau}|x_{\tau}) | x_t \right]$$

where $\lambda = 0.99999$ is the discount factor for an infinite time horizon. This particular discount factor means that population utility 100 years hence will still retain 99.9% of its current value, in keeping with the desire to protect exploited resources for use by future generations (Sumaila and Walters 2005). Population utility for the Pink-footed Goose is expressed as:

$$utility = \begin{cases} 0 & \text{if } N \leq 53.7 \text{ or } N \geq 90.1 \\ f(N) & \text{otherwise} \end{cases}$$

where $f(N) = -14.257 + 0.424N - 0.003N^2$ and N is the expected population size (in thousands) in the subsequent spring.

Thus, the objective function devalues harvest quotas that are expected to result in a subsequent population size different than the population target, with the degree of devaluation increasing as the difference between population size and the target increases. This current utility function, based on the updated population target of 72,000 birds, differs slightly from that used in previous years in that it utilizes consensus-convergence values from the formal balloting of stakeholder satisfaction in different values above/below the target (Madsen et al. 2025a). To account for this, we fit a quadratic curve reflecting how satisfaction with population size reaches a peak at a preferred target and declines as population size either falls below or above the preferred target, rather than across a range of values deemed ‘satisfactory’ close to the previous target of 60,000 (Figure 3.1-2). The optimal harvest strategy was computed using the publicly available software MDPSolve (© 2010 – 2011 Paul L. Fackler, <https://github.com/PaulFackler/MDPSolve>), which is a set of SDP tools written in the proprietary MATLAB® programming language.




Fig. 3.1-2. Utility functions used to derive optimal harvest strategies for Pink-footed Goose, each expressing the relative satisfaction with varying population sizes and both normalized to values from 0 to 1. The “old” function relates to the former spring population target of 60,000 geese, whereas the “new” function relates to the agreed spring population target of 72,000 geese (see Madsen et al. 2025a for details). The vertical red line represents the proposed Favourable Reference Population (FRP) size which has been set at 49,000 individuals.

3.1.5 Population status

a) Abundance and trends

As described in previous reports (Johnson et al. 2024, Sørensen et al. 2025), we have encountered some difficulties with count biases in the spring and autumn surveys, and the IPM has recently had difficulty interpreting the decreasing May count and the increasing November count. In 2023, the Data Centre committed to investigate the magnitude of bias in the biannual counts by exploring the use of GPS-tagged birds to estimate detection probabilities. A total of 109 Pink-footed Geese (mainly adult females) were GPS-tagged in western Finland in spring 2017 and 2018, in Svalbard in 2018, in Denmark in 2021 and in Norway in 2022 (Madsen et al. 2023; Schreven et al. 2024). However, precise positions of counted birds have only been available since November 2022.

We compared the time and location of counts with that of GPS-tagged individuals in the November 2022-2024, and May 2023-2026 censuses. In autumn 2025, we were not able to attain the GPS positions of tagged individuals because signals were jammed, probably due to military-induced interference with global navigation satellite systems (Jiguet et al. 2025). We tallied the number of GPS tags not present in a counting area on the day of the count, those recorded once in a counting area, and those recorded twice (i.e., double counted). We pooled GPS tallies from the November 2022, November 2023, and November 2024 censuses to increase sample size. We did not, however, pool all GPS tallies from the four May censuses. In May 2024, 2025, and 2026, the locations of GPS-tagged individuals at the start of the counts were provided to the observers in Trøndelag, Norway (where the majority of the population is concentrated in May) to increase the probability of a complete census, whereas these locations were not provided in May 2023. Observers provided GPS-positions of their counting points and time of visit. Based on the counts of GPS-tagged birds not present, counted once, and counted twice in a circle of 1 km radius around the counting points and within +/- 1 hour around the visit by the observer, the estimated detection probability during the May 2023 count was 0.82 (sd = 0.08), indicating a negative bias. In May 2024-2026, however, the pooled estimated detection rate was 1.03 (sd = 0.05), indicating a positive bias (Figure 3.1-3). During the November counts the estimated detection probability pooled across the three years was 0.84 (sd = 0.04), indicating a negative bias (Figure 3.1-4). Detailed methods for the comparison of counts versus GPS-tags are available from the EGMP Data Centre on request. For all November counts as well as May counts prior to 2024, we assumed the estimated means and sampling variances were constant over time but allowed year-specific detection probabilities to be drawn from these distributions. For the May 2024, 2025, and 2026 counts, detection probabilities were drawn from the pooled distribution of detection rates for these three years (Figure 3.1-3).

Accounting for these GPS-based biases resulted in little change in population estimates prior to about 2017, partly because estimates of spring population size based on a capture-mark-recapture program (since discontinued) were able to “arbitrate” between the May and November counts (which are both biased) up until this point. The greatest difference in population estimates based on the new GPS corrections occurs in the most recent years, which coincides with when the IPM was previously having trouble reconciling the difference in raw counts between May and November of the same year (as was originally reported in 2023).




Fig. 3.1-3. Estimated distributions of detection rates during May counts of Svalbard Pink-footed Geese based on the co-location of counted geese and GPS-tagged individuals. Values below 1.0 indicate a negative bias in the counts, while values greater than 1.0 indicate a positive bias (i.e., double counting). K represents the number of GPS tags active (included in counts) each season. For the current version of the IPM, detection rates for May counts prior to 2024 were drawn from the May 2023 distribution, representing the estimated negative bias in those years. May 2024, 2025, and 2026 counts were corrected for their positive bias (where locations of GPS-tagged birds were provided to observers) based on the pooled distribution (grey).




Fig. 3.1-4. Estimated distributions of detection rates during November counts of Svalbard Pink-footed Geese based on the co-location of counted geese and GPS-tagged individuals. Values below 1.0 indicate a negative bias in the counts, while values greater than 1.0 indicate a positive bias (i.e., double counting). K represents the number of GPS tags active (included in counts) each season. For the current version of the IPM, detection rates for all November counts were drawn from the pooled distribution (grey).

Estimated population sizes in May and November are provided in Figure 3.1-5. Following relatively higher May abundances in the past few years leading to a November 2025 population estimate of 103,715 (85,569 – 123,544), the May population size for 2026 appears to have declined slightly to an estimated 79,602 (60,374 – 102,021) birds, which is still above the target value of 72,000.




Figure 3.1-5. IPM-based estimates of abundance of Svalbard Pink-footed Geese in May and November, relative to the target of 72,000 birds. 95% credible intervals are indicated by the dashed lines.

b) Mortality and trends

Posterior estimates of country-specific harvests of Svalbard Pink-footed Geese are provided in Figure 3.1-6. Posterior estimates of annual harvest and survival rates of the flyway population are provided in Figure 3.1-7. Harvests and harvest rates were increasing prior to the implementation of the adaptive harvest management program in 2013 but have been somewhat stable since. We note that harvest has decreased substantially in Denmark during the last few years for reasons that are unclear. Probably it is related to geese using non-traditional areas and because the Danish Hunters' Association – following the decision taken during IWG8 – encouraged Danish hunters to reduce their offtake in 2023/24 due to uncertainty about population size. However, Danish harvest numbers for 2025 were unavailable at the time of writing. Estimates of annual survival have generally decreased during the entire period of record, although there is quite a bit of uncertainty associated with the estimates in the last few years (due to the cessation of the capture-mark-recapture program). In particular, the apparently large increase in annual survival in 2023 and then sharp decline in 2024 and 2025 should be viewed with some caution.




Figure 3.1-6. IPM-based estimates of harvests of Svalbard Pink-footed Geese (95% credible intervals are indicated by the dashed lines).




Figure 3.1-7. IPM-based estimates of harvest and annual survival rates of adult Svalbard Pink-footed Geese (95% credible intervals are indicated by the shaded polygons).

c) Reproduction and trends

Estimates of productivity, as indicated by the post-breeding proportion of young in the population, have been variable, with an average proportion of 0.19 ($se = 0.01$) young (Figure 3.1-8). Productivity has generally increased over the period of record and is highly correlated with the increasing number of days in which the mean air temperature is above freezing in May in Svalbard. The post-breeding proportion of young reached a maximum of 0.37 (0.32 – 0.41) in 2018 following a record 27 days above freezing in May in Svalbard. In contrast, the record low proportion of 0.13 (0.12 – 0.14) occurred in 1998, following 0 days above freezing in May in Svalbard. In 2025, the estimated post-breeding proportion of young was 0.24 (0.23 – 0.25), following 15 days above freezing in May in Svalbard. It should be noted that so far, the productivity reported here includes both the Svalbard and the Novaya Zemlya group (the latter only representing c. 4 % of the entire autumn population).




Figure 3.1-8. IPM-based estimates of the post-breeding proportion of young for Svalbard Pink-footed Geese (95% credible intervals are indicated by shaded polygon). In blue are the number of days above freezing in May in Svalbard.

3.1.6 Management guidance

The optimal harvest management strategy based on results of the IPM, candidate harvest quotas, and the objective function expressing the level of satisfaction with various population sizes (Madsen et al. 2025a) recommends harvest quotas ranging from 0 to a maximum of 20,000 (capped based on limited harvest capacity, reflecting a maximum harvest of 17,000 achieved in the past) within the most desirable range of population sizes around the target of 72,000 birds (Figure 3.1-9). Harvest quotas for population sizes <60,000 are very low unless the number of days above freezing in May in Svalbard is very high. Harvest quotas for population sizes >80,000 increase rapidly with small increases in population size, regardless of the number of days above freezing in May. For a population at its goal of 72,000, and with a mean number of 17 days above freezing, the harvest quota is ~16,000. Moreover, for a population near its target of 72,000, small changes in population size or days above freezing in Svalbard can lead to changes in quotas that are well below those which can be regulated effectively. The management strategy in Figure 3.1.9 also depicts the evolution of May population size, days above freezing in May, and harvest quotas since implementation of AHM in 2013.




Figure 3.1-9. Optimal harvest quotas for Svalbard Pink-footed Geese based on an IPM and an objective to maintain population size near 72,000. Thaw days represents the number of days above freezing in May in Svalbard. The black vertical dashed lines depict near-complete stakeholder satisfaction with population sizes. Also depicted are population sizes and days above freezing for the years in which AHM has been in place, with 13 = year 2013 and 26 = year 2026.

For 2026, the estimated breeding population is 79,602 individuals. May 2026 was relatively warm in Svalbard, with 19 days above freezing, and the predicted proportion of young in the autumn is 24%, which is well above average. This suggests a post-breeding population size of 98,706 individuals. Accordingly, the optimal harvest quota for the 2026/2027 hunting season is at the maximum capacity of 20,000 which, if achieved, would result in a spring population of ~78,000 in 2027. For comparison, the realized harvest averaged 9,903 (sd = 824) during the last three years of available data. If we use a quota of 20,000 and the agreed upon allocation of the quota (30% for Norway, 70% for Denmark), harvest quotas for Norway and Denmark this year are 6,000 and 14,000, respectively. During the last three years of available data for each, the harvest in Norway and Denmark averaged 2,925 (sd = 356) and 6,667 (sd = 361), respectively. Thus, harvests will have to be increased considerably to achieve the population target of 72,000 in spring. If harvest rates continue to be below 10%, we could expect further population growth.

3.2 Taiga Bean Goose *Anser fabalis fabalis*

3.2.1 Range states and populations

This chapter provides monitoring and assessment information for four populations of Taiga Bean Geese (formerly referred to as the Western, Central, Eastern 1 and Eastern 2 Management Units (MUs), respectively). The four recognised populations of Taiga Bean Geese (Figure 3.2-1) are delineated as:

- **Scandinavia (br) population:** Breeding in Northern and Central Sweden and Southern and Central Norway, wintering in Northern Denmark and Northern and Eastern United Kingdom;
- **Finland & North-western Russia (br) population:** Breeding in Northernmost Sweden, Northern Norway, Northern and Central Finland and adjacent North-western parts of Russia, wintering mostly in Southern Sweden and South-east Denmark;
- **Germany & Poland (nbr) population:** Breeding in upper Pechora region and western parts of west Siberian lowlands of Russia, wintering mostly in North-east Germany and North-west Poland.
- **Central Asia (nbr) population:** Breeding in the forested zone of Western and Central Siberia, wintering in China, Kazakhstan, and Kyrgyzstan.




Figure 3.2-1. Breeding and wintering areas of the four populations of Taiga Bean Goose. Lines indicate movement areas between breeding and wintering ranges.

In addition to the range states mentioned above, Taiga Bean Geese also occur regularly in Estonia, Latvia, Lithuania, Ukraine and Belarus during migration or in small numbers in winter. In The Netherlands, it has meanwhile become a vagrant species.

3.2.2 Population targets

The International Single Species Action Plan for the Conservation of the Taiga Bean Goose (TBG ISSAP) was revised and approved in 2025, with a planned lifespan of 12 years (2026-2037). In the revised plan, the agreed goal was to restore and/or maintain each of the four populations of Taiga Bean Goose to/in a Favourable Conservation Status (Alhainen et al. 2025). Using a reference-based approach, the Favourable Reference Population (FRP) for the Finland & NW Russia (br) population is set at the historic level of 57,000 individuals in

spring (Madsen et al. 2025b). Although sufficient data and capacity were only available to define a FRP for this one population, the revised TBG ISSAP aims to increase the population size of the Scandinavia (br) population to at least 2,000 individuals (spring), maintain the population size of the Finland & NW Russia (br) population above 70,000 individuals (spring), and stabilize the population sizes of the Germany & Poland (nbr) and Central Asia (nbr) populations with the hope of enabling their recovery and increase (Alhainen et al. 2025).

3.2.3 Management strategies

The size of the Scandinavia (br) population is currently considered too small to support hunting and the population is protected from hunting in the UK, Norway and in Denmark by a regional hunting ban. Taiga Bean Geese from the Germany & Poland (nbr) population are hunted in Belarus, Latvia, Russia and Poland, but the bag sizes in these range states are generally not known and data are insufficient to develop a sustainable harvest strategy. An effective protection of the wintering population of Taiga Bean Goose is in place in Germany, as all hunting on Taiga Bean Geese has been banned in the Federal State of Mecklenburg-Vorpommern. For the Finland & NW Russia population, the EGMP is operating under an interim harvest strategy intended to maintain the population size above 70,000 individuals (spring), while still providing limited hunting opportunities in Finland, Sweden and Denmark.

3.2.4 Assessment protocol

An annual stock assessment for the Finland & NW Russia (br) population is conducted by updating an Integrated Population Model (IPM), which was first adopted in 2020 and then revised in 2021 to exclude relatively small numbers of Tundra Bean Goose (*A. f. rossicus*) from count and harvest data. The IPM relies on harvest estimates (FI, SE, DK), and population counts in March (SE), October (SE), and January (SE, DK), along with mildly informative prior distributions for key demographic rates (a full description of the model can be found here: [TBG IPM](#)). The anniversary date of the IPM is March, with population size also estimated in the following months of October and January. The IPM predicts changes in abundance using a discrete, theta-logistic model:

$$N_{(t+1)}^M = N_t^M \left[(\psi(1 + \gamma_t) - 1) \left(1 - \left(\frac{N_t^M}{K} \right)^\theta \right) \right] - H_t$$

where N^M is the March population size, ψ is intrinsic survival from natural causes, γ is the intrinsic rate of reproduction, K is carrying capacity in the breeding season, θ is a parameter describing the type of density dependence (i.e., concave, linear, or convex), H is total harvest, and t is year.

Abundance in October, N^O , is predicted as a function of March abundance:

$$N_t^O = N_t^M + N_t^M \left[(\psi^{7/12}(1 + \gamma_t) - 1) \left(1 - \left(\frac{N_t^M}{K} \right)^\theta \right) \right] - H_t^F$$

in which we assume seven months of natural mortality, all reproduction and a portion of the total harvest occurring prior to October, where H^F represents the harvest in Finland.

Abundance in the following January is conditional on October abundance:

$$N_t^J = (N_t^O - H_t^D - \alpha H_t^S) \psi^{3/12}$$

where H^D and H^S represent harvests in Denmark and Sweden, respectively, and where α represents the proportion of the Swedish offtake occurring prior to January.

Abundance in the following March is thus:

$$N_{t+1}^M = (N_t^J - (1 - \alpha)H_t^S) \psi^{2/12}$$

where $(1 - \alpha)$ represents the proportion of the Swedish harvest that is taken after the regular season to help prevent crop damage (i.e., “conditional hunting”).

The IPM has generally been very successful at mediating all sources of count and harvest data to provide biologically reasonable and robust estimates of population status. This is encouraging because each data set has some degree of bias, including less than complete counts, missing counts from important parts of the range and a mix of Taiga and Tundra Bean Geese in both counts and harvest. However, over the last few years the IPM has had increasing difficulty harmonizing the data sources and providing reasonable results. With the addition of the 2023-2025 data, the IPM was unable to fully reconcile the various sources of data and so no reliable population assessment could be produced. It was suggested that this was attributable to changes in count methodology or observers, as well as unknown changes in the spatial and temporal distribution of birds, such that the extent of bias has likely not been relatively constant (as we have largely assumed). The full list of known problems with the count data, which had become worse over time, included:

- The March Bean Goose count in FI, SE and DK is critical as it is largely composed of Taiga Bean Geese and because it helps measure population status against the median recovery target of 70,000. The count is partly based on citizen-science data. Based on the observers’ advice, we have assumed a temporally constant number of 4,000 Tundra Bean Geese occur in that survey, which is probably not entirely accurate. Moreover, GPS tags available in March 2020 and 2021, suggested that perhaps 20% of Taiga Bean Geese occur outside the counting areas. But sample sizes were small, and we have no sense of whether the count bias is relatively constant over time.
- Until recently, the October count in SE has been higher than the March count, reflecting successful reproduction. But in 2021-2025, the October count has been lower than the March count, which is not biologically feasible unless there is a complete reproductive failure. Both citizen-science data from Finland in October in recent years and a recent analysis of detection rates during census periods (Piironen et al. 2023) suggests there may be large numbers of Taiga Bean Geese remaining in Finland at that time. We had heretofore assumed that Finnish birds had fully moved into SE prior to the count. Finally, GPS data from October 2019 suggest that a relatively large proportion of Taiga Bean Geese occurred outside the counting areas in SE, although it is important to note that sample size was very low and that it was unknown whether this count bias might change from one survey to another.
- Finally, there are the January Bean Goose counts in SE and DK. Counts in DE have not been made available to us, so we always knew the January counts were negatively biased. However, if the abundance in DE during January was relatively stable, then the January counts could still be useful in understanding population development. Unfortunately, the number of Taiga Bean Geese wintering in Germany has likely declined over time (Thomas Heinicke, pers. communication), perhaps due to birds remaining in more northern areas in response to warmer winters.

Of these known problems, the one that could be directly addressed for the current assessment is an estimate of the proportion of birds remaining in Finland at the time of the October count. Beginning in 2015 (and expanding drastically in 2019), colleagues at the Natural Resources Institute Finland (LUKE) began tracking Taiga Bean Geese with GPS tags, which has been supplemented with additional tags deployed in Sweden and Denmark (Antti Piironen, personal communication). From these tracking data, we were able to assess the proportion of GPS tagged birds present in Sweden (or within 50km of the Swedish border) at the time of the counts on October 15 of each year. Table 3.2-1 provides a breakdown of these data.

Table 3.2-1. The cumulative number of Taiga Bean Geese marked with GPS tags in Finland, Sweden and Denmark starting in 2015, and both the raw number of tagged birds and the total percentage of tagged birds occurring either in Sweden or within 50km of the Swedish border at the timing of the mid-October count in each year.

YEAR	# TAGGED BIRDS	# SWEDEN	PERCENT SWEDEN
2015	2	2	100%
2016	7	7	100%
2017	6	6	100%
2018	7	7	100%
2019	20	20	100%
2020	48	41	85%
2021	32	23	72%
2022	22	8	36%
2023	39	29	74%
2024	56	27	48%
2025	64	30	47%

From these data we can see a shift in the proportion of tagged birds occurring in Sweden beginning in 2020, which coincides with when October counts began to fall below the March counts. Since this time, the mean proportion of birds within Sweden on 15 October = 0.61 (± 0.19 SD), or 61%. From these parameters, we created a beta distribution (alpha = 3.28, beta = 2.14, n = 1000 samples) and randomly drew a correction factor for October counts in each year beginning in 2020. We repeated this process 1000 times, taking the mean correction value in each year and multiplied the October count by (1/mean correction value) to produce an updated count value for October in each year from 2020-2025.

3.2.5 Status – Scandinavia (br) population

a) Abundance

The size of the Scandinavia/Denmark and UK population of Taiga Bean Goose is assessed primarily at the wintering grounds in Denmark and Scotland.

In Denmark, a total of 710 Taiga Bean Geese were counted near Hundborg on 19 October 2025. This flock was assumed to include two flocks recorded in Thy prior to that: On 13 October, 51 birds had been recorded at Nors Sø, and 204 at Sjørring Sø/Rosvang in Thy. No Tundra Bean Geese were reported from Northwest Jutland in October. A dedicated Bean Goose count was carried out in Denmark a few days before the International Waterbird Census (IWC). Thy and the Pandrup area, the two known sites for birds from the Scandinavia (br) population, were visited on 11 and 13 January, respectively. The count resulted in a total of only 255 Taiga Bean Geese, all located in the central part of Thy. Another flock of 360 Taiga Bean Geese was found in the northernmost part of Thy on 18 January and assumed to have been over-looked during the first count. Hence, a total of 585 birds were observed in January 2026. Neither Tundra Bean Geese nor unidentified Bean Geese were reported from Northwest Jutland. While the October count was higher than in 2024, the January total was lower than in 2025 (Figure 3.3-2).

In Scotland, the wintering population is also relatively small, yet the behaviour of flocks utilising several different roosting and feeding areas during any given day makes it difficult to count the birds in the Slamannan Plateau landscape. This past winter, the highest number counted was 173 birds seen on 30 December 2025, with similar count results in mid-January (Brian Minshull, pers. comm.). Combined with the Danish January count, the highest number recorded this winter was 758 (Figure 3.2-2).

A relatively large number of Taiga Bean Geese were reported from a range of sites in the UK and elsewhere (including Ireland, the Faroe Islands and Spain) during the autumn and winter, yet none were observed at the traditional Norfolk sites of the Scandinavia (br) population. These birds were thought unconnected to the Slamannan wintering population as this remained stable compared to previous years, with the sighting of a single colour-ringed individual ringed in Finland suggesting instead a Finland & NW Russia (br) population origin. Sightings were widespread and generally in remote locations that were only occasionally visited, so an estimate of the total number of individuals is not available (Brian Minshull, pers. comm.; see Jones 2026 for further details).




Figure 3.2-2. Development of the winter population size of the Scandinavia (br) population of Taiga Bean Goose since 2005/2006. The number of Tundra Bean Geese and unidentified Bean Geese are included for Denmark. The dashed black line represents the population target suggested by Alhainen et al. (2025).

b) Survival

No survival information is currently available for the Scandinavia/Denmark and UK population of Taiga Bean Goose, although perhaps information based on ringed birds could provide some insight. The population is protected from hunting, but it is known that Taiga Bean Geese are accidentally shot in northwest Jutland in DK.

c) Reproduction

No age counts were carried out for this population during the winter 2025/26. The proportion of juveniles recorded in Scotland during previous winters can be seen in Figure 3.2-3).




Figure 3.2-3. Annual proportion of juveniles in the Scandinavia (br) population of Taiga Bean Goose since 2005. No data is available for 2025/26.

3.2.6 Status – Finland & North-Western Russia (br) population

a) Abundance and trends

Bias-corrected posterior estimates of population size at three times of the year are depicted in Figure 3.3-4. In March 2026, the population estimate was 75,470 (67,280 – 84,183), which is above the target of at least 70,000 individuals in spring. Seasonal estimates of population size were combined into a single time series in Figure 3.2-5. Recall that biological year runs from approximately 1 March in calendar year t to the end of February in calendar year $t+1$. Thus, we expect the lowest population in March following hunting and natural mortality in the previous winter, the highest population in October after reproduction occurs, and an intermediate population size in January following the hunting season. Our correction for the October population size (see Table 3.2-1) appears to have stabilized the fall population numbers to reasonable and consistent estimates for the past several years. However, lower than normal counts from Sweden in January and all of Fennoscandia in March 2026 indicate a slight population decline, despite fairly consistent growth rates over the past few years (Figure 3.2-6). While we cannot yet determine if these lower counts represent a true decline or a continuation of the known issues with the count data listed above without additional data, we draw attention to this potential decline both to note that management plans should exercise a degree of caution and to highlight the need for updated data collection protocols. Namely:

- Anecdotal reports suggest that Bean Geese may be migrating earlier in spring than in previous years. Spring counts (currently undertaken in second-third week in March) may need to be carried out earlier to get a better representation of abundance before migration begins.
- Concurrently, the location of GPS-tagged birds in Finland, Sweden, and Denmark should be investigated (similarly to what has been done with the October count) to determine the proportion of tagged birds outside of the region at the time of the March census. This would require a coordinated and sustained collaboration with LUKE and other partners deploying tags to update the information on a yearly basis to ensure changes in the spatial and temporal distribution of birds are accounted for.




Figure 3.2-4. Posterior estimates of population size (in black, with 95% credible intervals in grey) based on an IPM for Taiga Bean Geese in the Finland & NW Russia (br) population of Taiga Bean Geese. The horizontal line at 70,000 in the left panel represents the (spring) population target.




Figure 3.2-5. Posterior estimates of seasonal (March, October, January) population size (in black, with 95% credible intervals in grey) based on an IPM for Taiga Bean Geese in the Finland & NW Russia (br) population. The vertical dashed lines represent March of each year. The horizontal line at 70,000 represents the population target.




Figure 3.2-6. Posterior estimates of the population growth rate between March and October (in black, with 95% credible intervals in grey) based on an IPM for Taiga Bean Geese in the Finland & NW Russia (br) population.

b) Mortality and trends

Posterior estimates of country-specific harvests of the Finland & NW Russia (br) population are provided in Figure 3.2-7. Due to hunting restrictions in all three range states, the total harvest has only averaged 525 birds (sd = 125) since 2021. Posterior estimates of annual harvest rates and apparent survival of the population are provided in Figure 3.2-8. Harvest rates declined dramatically following the Finnish harvest moratorium in 2014, and this decrease in harvest rate coincides with strong growth in the population up to 2025. Estimates of apparent survival have likewise increased markedly over time and have remained fairly stable from 2021-2024. Note that because harvest estimates for 2025 are only currently available for Finland (approx. 100 birds), the 2025 estimates for Sweden and Denmark are derived completely from the IPM and not grounded in actual harvest numbers for that year.




Figure 3.2-7. Estimated harvests (in thousands, with 95% credible intervals in grey) based on an IPM for Taiga Bean Geese in the Finland & NW Russia (br) population. The vertical, dashed lines for Sweden and Denmark indicate the last year of data, where subsequent estimates were derived from the IPM (see text above for details).




Figure 3.2-8. Posterior estimates of harvest and apparent survival rate based on an IPM for Taiga Bean Geese in the Finland & NW Russia (br) population, with 95% credible intervals in grey. The vertical, dashed lines indicate the last year of data, where subsequent estimates were derived from the IPM (see text above for details).

c) *Reproduction and trends*

Posterior estimates of the intrinsic reproductive rate (i.e., absent any density-dependent effects that may have been operative) have varied little over the timeframe of the IPM (Figure 3.2-9) and have averaged 0.32 ($sd = 0.03$) (or approximately 24% young absent any density-dependent effects). It should be mentioned that posterior estimates are similar to their informative prior mean, albeit more precise.




Figure 3.2-9. Posterior estimates of the intrinsic reproductive rate (in black, with 95% credible intervals in grey) based on an IPM for Taiga Bean Geese in the Finland & NW Russia (br) population.

3.2.7 Status – Germany & Poland (nbr) population

As the two Russian breeding populations overlap, the size of the Germany & Poland (nbr) population is estimated by counts in the wintering areas. Although available information is still limited, the latest counts from northeast Germany in January 2020 and 2021 (Heinicke et al. in prep.) revealed a population size of 10,000-15,000 birds. The current wintering population in northern Poland is unknown, but recent counts have resulted in less than 500 birds counted in January, all found in Pomerania (Wylegała & Smyk, 2025), which is a much lower number than the 5,000 wintering birds previously estimated (Alhainen et al. 2025). Thus, the total population size of the Germany & Poland (nbr) population is estimated at 10,000-20,000 birds.

Taiga Bean Goose is officially protected in parts of their Russian breeding range (Volkov & Rozenfeld 2025). However, less than 16% of key sites are covered by the existing network of protected areas, and Taiga Bean Geese probably spend more than 80% of the migration period in unprotected areas (Rozenfeld et al. 2024, Volkov & Rozenfeld 2025). Based on the movements of 43 geese tagged with GPS/GSM transmitters during 2018–2023, almost 60% of all stopover sites are located in Russia (Rozenfeld et al. 2024, Volkov & Rozenfeld 2025).

3.2.8 Status – Central Asia (nbr) population

Obtaining population estimates to support accurate population estimates and trends for this population is difficult (Alhainen et al. 2025). As the breeding areas likely overlap with the Germany & Poland (nbr) population, estimates for the Central Asia (nbr) population should be based on wintering numbers in Central Asia, yet information from Kazakhstan, Kyrgyzstan, and China is limited. Thus, observations from Russia provide the most recent data.

Taiga Bean Goose numbers in Russia during the spring and summer (breeding and moulting), including both Russian breeding populations, are estimated at maximum 19,000 individuals (Red Book of the Russian Federation, 2021). In Kyrgyzstan, 48 Taiga Bean Geese were observed wintering at Lake Issyk-Kul in 2005, compared to records of almost 600 birds wintering in 1998 (Heinicke, 2009). Expert reviews of observations submitted to citizen science platforms during 2012-2024 indicate the presence of up to at least 500 individuals on the wintering grounds in China, Kazakhstan, and Kyrgyzstan (Berlioux, J., 2025).

No recent information is available on the survival and productivity of this population.

3.2.9 Management guidance

Birds in the Scandinavia (br) population are protected from hunting in all Range States yet may occasionally be exposed to hunting as the range of this population overlaps with those of several huntable goose species. Given the endangered status of this population, harvest restrictions should remain in place, and the extent of unintended and illegal offtake should be investigated. Furthermore, as the population status and population dynamics are poorly understood, improved monitoring and research efforts are encouraged in all Range States.

For the Finland & NW Russia (br) population, we suggest a continuation of the restricted hunting levels to ensure that the population stays above the target of 70,000 birds in spring, while continued investigation into the causes (biological or methodological) that have led to decreased counts in 2026 are undertaken.

Since the protected Germany & Poland (nbr) and Central Asia (nbr) populations are both small and probably declining in the long term, with fragmented ranges and possibly low survival, harvest restrictions should be maintained and enforcement strengthened. Improved monitoring in the wintering areas should be encouraged.

3.3 Greylag Goose *Anser anser*

This chapter compiles monitoring data on the population status of the NW Europe (br) population of Greylag Goose and provides an update on the establishment of the monitoring and modelling frameworks necessary to perform a dynamic and model-based assessment at MU level (Nagy et al. 2021c).

3.3.1 Range States and Management Units

The range states for the NW Europe (br) population of Greylag Goose include Norway, Sweden, Finland, Denmark, Germany, The Netherlands, Belgium, France and Spain. Geese belonging to this population also occur regularly in Poland, Czech Republic and Portugal, but these countries are not included as principal Range States as numbers recorded here constitute less than 1% of the total population. Based on the recognition of regional differences in migratory behaviour and the human-wildlife conflicts related to this population, it has been agreed to define two MUs (Nagy et al. 2021c).

MU1 includes the breeding populations in Norway, Sweden, Finland, and Denmark that subsequently stage and winter also in The Netherlands, Germany, and Belgium. Some birds from this MU migrate to the southernmost wintering sites in France and Spain. MU2 is the mainly sedentary populations of The Netherlands, Belgium, and Germany, including also a small French breeding population of c. 2400 pairs (estimated in 2021/2022) (Figure 3.3-1). Although the German population is generally regarded as sedentary, breeding birds in the eastern part of the country are known to show migratory behaviour (Bairlein et al. 2014).




Figure 3.3-1. Annual distribution and main migration routes for the NW Europe (br) population of Greylag Goose including breeding (medium grey) and wintering (light grey) areas, as well as areas which are both used during the breeding and wintering period (dark grey) as presented in the [ISSMP](#) (up for evaluation in 2030). The two management units (MUs) are also shown: MU1 for the migratory population (in green) and MU2 for the sedentary population (in blue).

3.3.2 Population FRPs and targets

The FRP for the breeding season is 31,100 pairs for MU1, 72,980 pairs for MU2 and 104,080 pairs for the whole population. The wintering FRP is 370,400 individuals for the entire population (Nagy et al. 2021). Targets for MU1 and MU2 are 70,000 and 80,000 breeding pairs, respectively, resulting in an approximate wintering population size of 545,000 individuals (Nagy et al. 2021c).

3.3.3 Management strategies

In the face of deep uncertainty related to estimates of population size and offtake at the flyway level, an information-gap (“info-gap”) decision model was developed in 2020 to allow decision makers to make informed choices about the magnitude of offtake until a dynamic, model-based management of the population could be established based on more reliable monitoring information (Nagy et al. 2021c; Johnson and Koffijberg 2021). As agreed, the info-gap decision model was ceased after a 3-year period, however the dynamic, model-based management has not yet been established as data on population size and/or offtake apparently continue to be biased (and, for some Range States, data availability is limited).

Based on the info-gap decision model, range states agreed on a management criterion of a 15% reduction in the flyway population size over 10 years, which meant an annual finite growth rate of 0.96 – 1.00 ([EGM IWG5 meeting report](#)). To move beyond the rather crude info-gap approach, the [AFMP](#) mandated the establishment of “an internationally coordinated population management programme for both [management units], including offtake under hunting and, if necessary, under derogations, encompassing monitoring, assessment and decision-making protocols” (Nagy et al. 2021c). Considerable progress has been made in this effort, including the development of a flyway population model, which characterizes the dynamics of both breeding segments (MU1 and MU2) and accounts for the mixing of the two segments during autumn and winter. Based on input from the IWG, a utility model for Greylag Geese has also been developed that describes the relative level of satisfaction among stakeholders as the number of breeding pairs deviate from their agreed-upon targets. This utility model can be used to evaluate various offtake strategies in terms of their ability to meet population targets.

It should be noted that the current modelling framework is used to simulate how varying levels of offtake in different seasons and areas might affect whether the MU populations are near their targets when the ISSMP comes up for review in 2030. It is *not* intended to prescribe the magnitude and distribution of offtake at this time because current estimates of offtake are apparently biased high. Moreover, we note that while derogation is a legal means of alleviating local socio-economic conflicts, it cannot be used in a planned manner to meet a population target. However, once more reliable empirical estimates of offtake are available, the model can be parameterised and used to forecast the population trajectory under those levels of offtake to help determine whether the population is trending toward the target or FRP (e.g., as is done with Barnacle Geese). Also, given reliable estimates of derogations, the model could be used to help prescribe the level and distribution of recreational hunting to help attain population targets.

3.3.4 Assessment protocol

a) Population model

We use a post-breeding projection matrix, decomposed into summer and winter components. The summer component consists of the two breeding management units (MU1 and MU2), and the winter component consists of two wintering areas (North and South) (Figure 3.3-2). There is a broad overlap in the wintering distributions of the two breeding units. The southern unit is largely comprised of MU1 birds and is of special concern as short-stopping of migratory birds may eventually cause the range to fall below the FRR.

We also divide the annual cycle of Greylag Geese into a breeding season (March – August) and a wintering season (September – February) (Figure 3.3-3). We recognize the definition of seasons is somewhat arbitrary as it must represent a compromise of phenology that varies among countries.




Figure 3.3-2. Breeding management units and wintering areas for the NW Europe (br) population of Greylag Goose.




Figure 3.3-3. Diagrammatic representation of the model for the annual cycle of the NW Europe (br) population of Greylag Goose.

The model was parameterized using basic life history information and some limited empirical data (Appendix A.3). The model can be improved with a time-series of post-breeding population sizes in each MU, with the proportion of young in those counts, seasonal (March – August, September – February) offtake by country, and winter counts by country. The summer age ratios are particularly important in helping determine the number of breeding pairs, which is the criteria used in the MU-specific population targets. The biggest obstacle to model improvement and application, however, continues to be the acquisition of reliable empirical estimates of seasonal offtake.

b) Utility function

The effort to better coordinate the offtake of Greylag Geese involves specifying objectives and their relative importance in managing the abundance of Greylag Geese. Beyond an objective to maintain the population in a favourable conservation status, the objectives specified by the [ISSMP](#) are depicted in Figure 3.3-4. However, the [ISSMP](#) did not prioritize these objectives, and so the IWG was asked to specify their relative importance (also shown in Figure 3.3-4). These objectives and their weights were used to specify population targets of 70 and 80 thousand breeding pairs for MU1 and MU2, respectively (Johnson et al. 2021).




Figure 3.3-4. Relative importance of seven objectives for managing the offtake of the NW Europe (br) population of Greylag Geese. Management objectives are to maximize cultural and aesthetic values (cav), minimize agricultural damage (crop), minimize management costs to governments (cost), minimize deleterious impacts to habitats (hab), maximize satisfaction with the level of recreational hunting (hunt), minimize amenity fouling and disease transmission (health), and minimize bird strikes to aircraft (strike). From Johnson et al. 2021.

3.3.5 Population status

a) Abundance

The population size of the NW Europe (br) population of Greylag Goose is assessed twice a year, during winter and more recently during the post-breeding period in summer/early autumn. The winter abundance represents the total flyway population size, and the post-breeding abundance represents the size of each management unit.

Winter abundance is routinely estimated using the International Waterbird Census (IWC), as well as values from special goose counts in Denmark and the Netherlands. Based on those sources, the estimated population size was 802,320 individuals in January 2025 (Figure 3.3-5), similar to the result of the winter population count in 2023, which indicates a decrease in population size of around 130,000 individuals compared to January 2024. However, as some Range States have yet to report the final IWC results from 2025, the population estimate may not accurately reflect population development, and it remains to be seen whether the population growth has levelled off or will continue in the coming years. As mentioned in Heldbjerg et al. (2021), estimates from Spain included

a high degree of imputing due to data gaps during 2010-2013, which may have resulted in an overestimation of the actual population size by some 200,000 birds during those years. In recent years, Spanish data are not considered to be severely biased, yet we have chosen to keep both graphs (including and excluding Spain) in Figure 3.3-5. The imputed IWC value for January 2025 indicates that around 6,000 Greylag Geese wintered in Spain this year, potentially reflecting continued decline in recent years.




Figure 3.3-5. Development of the size (number of individuals) of the NW Europe (br) mid-winter population of Greylag Goose based on IWC imputed values from 1980-2025, including (solid line) and excluding (dot-dashed line) estimates from Spain. The dashed black line represents the target for the wintering population, and the red dashed line represents the wintering FRP.

Post-breeding abundance estimates are achieved through a combination of long-running and recently established national initiatives. For MU1, annual post-breeding counts have been carried out in Denmark and Sweden during September for decades (Nielsen et al. 2023, Haas et al. 2023). In 2022, Denmark organized an August count to provide a better estimate for the national population size (Jensen et al. 2023a), after which the national September count was moved to August and will be carried out biennially from 2023 onwards. Birds in Finland have also been counted during summer in recent years, but as these birds are assumed to be included in the Swedish September count, they are currently not added to the annual total for MU1. In Norway, counts were carried out at selected sites in August 2022, and a model-based estimate for the population size was subsequently produced, including also previous counts at selected sites, data from the Norwegian breeding bird monitoring scheme, and national hunting bag statistics (see Sørensen et al. 2024). For MU2, counts are carried out and available from parts of Germany (Nordrhein-Westfalen, Niedersachsen and Schleswig-Holstein), The Netherlands and Belgium (Niedersächsische Sommer-Gänsezählung 2025, Koffijberg & Kowallik 2024, Wolff et al. 2023). Numbers from France and Spain are currently regarded as non-essential due to small breeding populations, in some cases originating from feral geese. The number of breeding pairs is estimated every six years in France.

Counts from the post-breeding period produced a minimum of 540,115 individuals in 2022 for MU1, with counts in Denmark and Sweden producing similar results in the following year(s). In 2025, a total of 447,947 individuals

were counted in Denmark (August) and Sweden (September) (Figure 3.3.-6). In MU2, 808,171 birds were reported in 2025 (no data available from France, and from Germany data are only available from Nordrhein-Westfalen, Schleswig-Holstein, and Niedersachsen, i.e., only three of the 16 Bundesländer) (Figure 3.3-7).

In 2024, we investigated how the number of breeding pairs in the spring might be back-calculated from post-breeding censuses for the two management units of the NW Europe (br) Population of Greylag Goose (see Sørensen et al. 2024 and further details provided in Johnson et al. 2024). Although our methods only provide a rough approximation for the number of breeding pairs because empirical data are insufficient to do otherwise, such calculations will help evaluate progress in terms of reaching the agreed targets of 70,000 pairs in MU1 and 80,000 pairs in MU2.

The most recent year of complete data is 2022. For MU1, for a post-breeding population of 540,115 (Sweden, Finland, Denmark, and Norway) the estimated number of breeding pairs was 132,142 (113,293 – 150,852). Around the middle of the last decade, the number of breeding pairs (all countries) in MU 1 was estimated at 84,000 (S. Nagy, personal communication). For MU2, we used a 2022 post-breeding population of 769,875 (Netherlands, Belgium, and Germany (Nordrhein-Westfalen, Niedersachsen, and Schleswig-Holstein)) and a spring population in France of 8,323, resulting in an estimate of 185,432 (147,605 – 206,367)² breeding pairs. Be aware that this was an underestimate of the total population size in MU2 as data was only available from three German federal states. Around the middle of the last decade, the number of breeding pairs (all countries) in MU2 was estimated at 139,400 (S. Nagy, personal communication). Since 2022, population estimates based on post-breeding counts have increased for all Range States conducting such counts (see figures 3.3-6 and 3.3-7).




Figure 3.3-6. Number of Greylag Geese counted during the post-breeding counts in MU1 Range States, consisting of available data from Denmark (September 2005-2021, August 2022-2025), Sweden (September 2005-2025), and Norway (2022). Birds breeding in Finland are assumed to be included in the Swedish count.

² Note that this estimate has been updated in 2026 based on revised summer counts from Germany.




Figure 3.3-7. Number of Greylag Geese counted during the post-breeding counts in MU2 Range States, consisting of available data from Belgium (2010-2018 and 2021-2025), the Netherlands (July 2005, 2009, 2012-2025), Germany (Nordrhein-Westfalen July 2011-2025, Niedersachsen July 2018-2025, Schleswig-Holstein June/September 2019, 2021-2024), and France (2021-2022).

b) Survival and mortality

1) Offtake at population level

Hunting bag estimates are available from all range states except France and sum to 176,297 for the 2024/2025 season. Derogation data from 2024 are available from all range states where derogations have taken place, although data is missing from one Dutch province, and indicate that 273,305 geese were killed under derogation (including lethal scaring permitted at municipality level in Norway, following Norwegian game legislation). A significant reduction in the number of birds killed under derogation has occurred in Denmark, possibly as a result of increased efforts by the national authorities to control that preconditions for granting derogation permits are met.

Data suggest a minimum offtake of 449,602 Greylag Geese in 2024/2025 (Figure 3.3-8), indicating that offtake remains relatively stable across the Range States. Given an estimated summer population of around 1.3 million birds in recent years, and an estimated winter population size of 802,320 individuals in January 2025 (see above), with no indication of a declining trend, there is however reason to believe that offtake data are still biased high. We will continue to investigate this bias further in the coming years, with support from the relevant range states.




Figure 3.3-8. Total number of Greylag Geese killed under derogation (per calendar year 2013-2024) and hunting (per season from 2013/2014-2024/2025). Data from the Netherlands are incomplete (missing from one province) for 2023 and 2024, and in some years data from Spain include only offtake in Andalusia. Derogation numbers from Norway are estimates, based on information from county governors and municipalities. Offtake in France during 2024/2025 is not yet available.

II) Survival

No updated monitoring results available.

III) Crippling

In contrast to the low crippling rate seen in Pink-footed Goose in recent years (Buij et al. 2026), the crippling rate of Greylag Geese appears to remain relatively high. In 2022, an overall crippling rate of 21 % was found among Greylag Geese caught in Sweden (Månsson et al. 2024), with regional results ranging from 11.7 % to 26.4 %. Updated results from Sweden (n=139) and the Netherlands (n=622) covering the period 2017-2022 have demonstrated adult crippling rates among Greylag Geese of 22 and 30%, respectively, in the two countries during summer (Buij et al. 2026). As study periods overlap, these results are not indicative of a trend. However, the crippling rates presented in Buij et al. (2026) are based on a much larger sample size, covering both MUs, thus indicating a potential cause for concern across the flyway.

c) Reproduction

In MU1, age counts have been carried out in two Range States in 2025. Thus, new information is available from one region in Norway (Vesterålen) and a range of sites in Sweden. In Vesterålen, juvenile percentages have ranged between 4.1 and 40.1% during the years 2020-2025, with the most recent estimate being 32.5%. In Sweden, the juvenile percentage reached a peak in 2024 at 25.0% (Haas et al. in prep.), indicating an unusually productive breeding season following a couple of years with juvenile proportions of 5.0-7.5% (Haas et al. 2023). In 2025,

the surveyed flocks contained 20.1% juveniles, thus overall productivity in MU1 appears to remain high (Figure 3.3-9).

For MU2, extensive age counts are available from the Netherlands (Hornman et al. 2024, Sovon Vogelonderzoek Nederland) and North Rhine Westphalia in Germany (Koffijberg & Kowallik 2024). Limited sampling in Niedersachsen (Germany) has also been carried out since 2019. After an initial peak in the Netherlands in the late 1990s, the proportion of juveniles declined markedly, resulting in an average juvenile proportion of 18.0% in the Netherlands and 19.3% in Germany during the last decade. However, peak years with more than 20% juveniles are seen in both Germany and the Netherlands (Figure 3.3-9).




Figure 3.3-9. Proportion of juveniles in the NW Europe (br) population of Greylag Goose in five Range States: Norway (Vesterålen 2020-2022 and 2024-2025, Oslofjord Area 2020-2024), Finland (2022-2023), the Netherlands (1997-2024), Germany (North-Rhine Westphalia 2011-2025, Niedersachsen 2019-2020 and 2022-2025), and Sweden (2022-2025).

3.3.6 Management guidance

In 2023, using the preliminary population model, we simulated all permutations of offtake rates of 0.00 – 0.40 in increments of 0.02 for all seasons and areas (194,481 offtake scenarios). We retained all offtake strategies that had a high probability of meeting both MU targets by the time the [ISSMP](#) is due for revision in 2030 (Figure 3.3-10).

The simulations of the preliminary model demonstrated that no unique level and distribution of offtake would meet MU population targets. Rather, alternative approaches to coordinating offtake must be evaluated ultimately not only in terms of their ability to meet population targets, but also in terms of cost, feasibility, and legal mandates. The 50 offtake strategies with high probability of meeting the MU targets are of two basic types: (a) those with relatively high spring/summer derogation and low winter offtake, and (b) those with low spring/summer derogation and relatively high winter offtake (Figure 3.3-10). In June 2023, the IWG recommended that offtake be concentrated during the wintering period to the extent possible. Again in June 2024

and 2025, the IWG agreed that all Range States will focus on offtake strategies that minimise the need for breeding season derogation.




Figure 3.3-10. Fifty alternative offtake strategies for Greylag Geese with high probability of meeting the MU targets by 2030, ordered by decreasing level of total offtake. Values of offtake are the means over the timeframe. Colours correspond to those used in Figures 3.3-2 and 3.3-3.

Comparing the mean levels of offtake for the two sets of management strategies (a and b) with current estimates of offtake implies either that the flyway population is underestimated or the estimates of offtake are biased high, perhaps extremely so. To reconcile discrepancies between reported levels of offtake and those needed to meet population targets, the following data are needed in descending order of priority:

1. Reliable offtake estimates: by country and biannual period (spring-summer: March-August and fall-winter: September-February) for the most recent five calendar years.
2. Summer or early autumn abundance: by country for those conducting such surveys; all years in which they are available.
3. Post-breeding age ratios: all years and countries where available; should include counts of young and total sample size. Data should be provided at the lowest level available (e.g., by flock or location).
4. Winter counts: all years and countries where available.

While data availability has greatly improved in recent years, we still have reason to suspect that either offtake or population counts are biased. While investigating ways to improve the management guidance for the NW Europe (br) population of Greylag Goose, emerging patterns in the migratory behaviour of the population have been identified, and the associated management implications are currently being investigated by an ad-hoc group established following EGM IWG10 (see document [AEWA/EGMIWG11.24](#)). Primarily, the work of the ad-hoc working group will improve MU-specific population size estimates and shed further light on the changing distribution of geese through increased short-stopping. This may affect the mean levels and the recommended distribution of offtake between seasons in each MU to reach the population target. At the same time, methods for reporting offtake in the Netherlands are currently undergoing a revision which may affect the overall offtake level.

For the reasons mentioned above, the EGMP Data Centre has postponed a review and revision of the existing population model until 2027, hoping that updated information on population size and offtake will be available for a thorough investigation of population dynamics. Our management recommendations for the coming season are thus to continue focusing on offtake strategies minimizing the need for spring/summer derogation. As the current level of offtake does not seem to jeopardize the population, we see no immediate need for adjustments to the national legislation. However, despite the continued growth in both MUs, we do wish to highlight the declining number of Greylag Geese from MU1 wintering in the southern part of the range and that these birds will be subject to an increasing harvest rate if the current level of hunting is maintained while numbers continue to decline.

3.4 Russia breeding population of Barnacle Goose *Branta leucopsis*

This assessment of the population status of the Russia (br) population aims to investigate the cumulative impact of derogation and hunting (where legally allowed) on the status of the flyway population and the status of the three individual management units (MUs) which have been defined. In addition, it provides guidance for management in 2026/27 and thereafter, for instance covering coordination of derogation measures among countries within one MU, in case the assessment calls for this. Besides, gaps in the monitoring networks are addressed. In line with the framework set out in the AFMP (Nagy et al. 2021b), the assessment is based on an Integrated Population Model (IPM). This model was initially developed for the Russian breeding population (MU1) only and presented during IWG5 in 2020 (Baveco et al. in Nagy et al. 2021b). In 2022, it was extended to the Baltic and North Sea breeding populations and then used in a first full assessment of the population status in 2005-2021 (Jensen et al. 2022). During IWG7 in 2022, it was decided to use the model framework of the IPM for an annual update making use of the newest available monitoring data. In autumn 2022, the IPM was further refined with input from a review made by the EGMP Data Centre and the Norwegian Institute for Nature Research (F. Johnson, K. Layton-Matthews). The current iteration of the IPM reflects the version that has been used after this review, for the assessments 2023-2025.

3.4.1 Range States and Management Units

The range states for the Russia (br) population of Barnacle Goose include Russia, Finland, Estonia, Sweden, Norway, Denmark, Germany, the Netherlands and Belgium. Among these range states, three management units have been delineated, covering the (Arctic) Russian breeding population (MU1, migratory and containing by far the largest share of the total flyway population), the Baltic breeding population in Finland, Sweden, Estonia, and Denmark (MU2, migratory), and the North Sea breeding population in Germany and the Netherlands (MU3, sedentary) (Figure 3.4-1). Formally, the Norwegian population in MU2 (now expanding from original breeding sites in the greater Oslofjord region) and the Belgian population in MU3 are not covered by the AFMP, as these populations are not recognized as naturally occurring, but derived from feral populations, by the respective country administrations. Still, these birds (altogether < 5,000 individuals and less than 1% of the flyway population) mix with the other birds in winter, so they are included in the monitoring setup and in the input data for the IPM.

During winter (i.e. from October to mid-May), birds from all management units mix in Sweden, Denmark, Germany, the Netherlands and Belgium. All these countries organize the traditional midwinter count (part of the International Waterbird Census) or dedicated goose counts, usually with a very good coverage of the wintering sites of Barnacle Geese. At present, the Netherlands and Germany are still the most important wintering countries, supporting on average about 47% and 27% of the flyway population, respectively (averages from census data collected in January 2023-25). However, in the past ten years other countries, especially Denmark, has received an increasing share of the wintering population while the wintering range has expanded into southern Sweden.

3.4.2 Population FRPs and targets

The FRPs for the breeding season have been defined as 113,000 pairs for MU1, 12,000 pairs for MU2 and 12,000 pairs for MU3 (Nagy et al. 2021b). The FRP for the entire population has been set at 380,000 individuals in winter, reflecting the flyway population size in 2000, when AEWA came into force (Nagy et al. 2021b). As Barnacle Goose is an Annex 1 species of the EU Birds Directive, the AFMP does not aim to maintain the population at or reach a pre-defined target level. Impacts of offtake under derogation is evaluated in retrospective, and management is carried out by each single EU Member State under the conditions for derogation, outlined in Art. 9 of the EU Birds Directive and implemented in national legislation of each EU Member State. Birds in Norway (not an EU Member State) have a similar protective status, following Annex II of the Bern Convention, including derogation-like measures (lethal scaring, not for regulating numbers) to prevent crop damage (in this case granted

by, and reported to, the municipalities), if other measures of prevention have failed. Hunting (harvest) is only carried out outside the EU, mainly in Russia.

3.4.3 Management strategies

The AFMP aims to prevent the population or any of its MUs from declining below the specified FRPs (Nagy et al. 2021b). Hence, the FRPs represent the lower limits of the legally acceptable population sizes, but as such do not reflect true targets for population size. Monitoring of the population size and offtake and predictive modelling (IPM) of the cumulative impact of national derogation measures and hunting (where it is legally allowed) is used to inform national decision-making during the IWG-meetings to ensure this. The cumulative impact of derogation and hunting on the development of the population is assessed periodically, along with the likelihood of serious damage to agriculture, risk to air safety and to other flora and fauna, and the non-lethal measures taken to prevent damage/risk, as well as the effectiveness of these. This means that once every year, additional monitoring data provide insight into the population development, whereas once every six years an evaluation of the AFMP is carried out, assessing the implementation progress through the pre-defined indicators (next assessment due in 2026). Within this framework, it has also been agreed to coordinate monitoring of the offtake under derogation and hunting when the size of the population size (of single MUs or for the entire population) is below 200% of the FRP, as a precautionary measure. This includes monitoring of population size, offtake, prediction of population development (by the IPM), and coordination of offtake and conservation measures when necessary. A protocol for this coordination has been subject to discussions in the Task Force for the Russian/Netherlands and Germany population of Barnacle Goose (see doc. AEWA/EGMIWG/7.14 from EGM IWG7 in 2022). So far, it has only been applied in MU3, to avoid the population falling below FRP. During IWG10 it was also discussed whether coordination is needed in MU2 (see also section on Management Guidance).




Figure 3.4-1. Management units of the Russia (br) population of Barnacle Goose.

3.4.4 Assessment protocol

The assessment of the status of the Russia (br) population is carried out using an IPM. Input for the model is derived from monitoring data on abundance, productivity, and offtake under derogation, both for summer and winter situations (see Appendix A.4 for coverage in each country and the [EGMP Database](#) for an overview of data used). The way the IPM framework accounts for the impact of offtake in the respective management units is shown in Figure 3.4-2. Monitoring data on abundance and productivity have been included up to January 2025 (winter flyway size, based on midwinter counts) and summer 2024 (numbers per MU, based on summer counts). For derogation data the last year taken into account was 2024 (submission of data on offtake under derogation from 2025 to the EU is due September 2026). In case of missing abundance data or incomplete time series, annual growth rates or estimates have been used to provide estimates. This was the case for Schleswig-Holstein in Germany in January 2025 (data not published yet) and for Sweden in January 2021 and 2022 (data missing due to very low coverage during the corona pandemic). Previously missing data from Germany has been provided through recently published census data.

Because summer counts are completely missing prior to 2005, results of the assessment shown in this chapter solely refer to the period 2005-2025. An overview of the longer time series is included in the [EGMP Database](#) and the annual status report from 2021 (Heldbjerg et al. 2021). This is especially relevant for productivity, which has declined in the Russian population in the long term, but less so when considered from 2005 onwards. As in the status report from 2025 (Sørensen et al. 2025), the results from September counts have been used in Sweden as a proxy for the size of the summer population. The count is carried out in mid-September, before migratory birds from MU1 have arrived (F. Haas, pers. communication), which has been confirmed by analyses of resightings of marked birds (L. de Vries, pers. communication). Moreover, exchange with the Finnish summer population is considered low, as the Finnish count is done only two weeks earlier than the Swedish count. Nevertheless, the exchange between Finland and Sweden (and perhaps also Denmark) is an issue which still needs further investigation, and preferably confirmation by transmitter- or ringing data (this is work in progress). A complete count in summer, covering all relevant parts of Sweden, is still not considered feasible, so using the September count instead is regarded as the best alternative option at present. Also, gaps in time series for Danish numbers during summer have been interpolated from periodical counts available. Currently, Denmark is the main gap in collection of census data during the summer period (see also Appendix A4).

Following a review by the EGMP in autumn 2022, the IPM was adapted in several ways. A simplification resulted from equating survival from natural mortality in the summer to that in the winter period. A reanalysis was performed of within-year variation in juvenile counts, and different approaches in defining the associated priors were tested. An approach for evaluating the goodness of fit, based on post-predictive checks, has been implemented as well, following the approach taken in the Pink-footed Goose IPM (Johnson et al. 2022).




Figure 3.4-2. Overview of the offtake of Barnacle Geese in the different regions experienced by the birds belonging to the different MUs of the Russia (br) population. Local breeding populations (green boxes) in the Baltic Sea and the North Sea areas experience offtake around the breeding period (“B” and “N” respectively). Outside this period (blue boxes), birds of all three MUs experience offtake in their staging and wintering areas (“BS” and “NS” respectively). The scheme is simplified, as in the model and data the first set is split in offtake before and after July 15, and the second in offtake before and after January 15. Half-yearly survival is effectuated directly before and after offtake in staging and wintering areas (“BS” and “NS”). Offtake in Russia is unknown. RU: Russia, FI: Finland, DK: Denmark, SE: Sweden, EE: Estonia, NL: Netherlands, DE: Germany.

3.4.5 Status

a) Abundance

Posterior estimates from the IPM indicate a flyway population size of about 1.5 million individuals (rounded) in January 2025, which is slightly below the numbers estimated for 2024 (Figure 3.4-3). Results from the IPM and results from field counts correspond very well, with all counts being within the 95% credibility intervals of the IPM-estimates. The IPM-estimates for January 2025 resemble those in January 2021-2022, suggesting that the total flyway population size has not undergone an overall growth in the past five years. On a longer term, however, there has been a clear increase (see Helldbjerg et al. 2021) but the rate of increase has levelled off in the past decade. The estimated population size in January 2025 is four times the FRP (100% and 200% levels shown by the dashed line in Figure 3.4-3). Census data indicate that in January 2025, especially wintering numbers in the Netherlands were higher than average (800,000 – second highest winter count recorded so far). Higher numbers in the Netherlands corresponded to lower numbers in Northern Germany (perhaps as a result of a short cold spell prior to the census period), while in Denmark and Sweden numbers were the highest recorded since the start of the counts. On a longer term, wintering numbers in the Netherlands have stabilized since 2012/13 (Hornman et al. 2024) while especially in Germany, Denmark and Sweden they have continued to grow until recently (Blüml & Kruckenberg 2023, Nielsen et al. 2024, Haas 2025), implicating a northeastward expansion of important winter concentrations in the past decade.

Based on the posterior abundance estimates for July, the Russian breeding population (MU1) is by far the largest of all MUs, amounting to approx. 1.6 million individuals (rounded) in July 2024, whereas the Baltic population in MU2 and North Sea population in MU3 are much smaller: 64,000 and 67,000 individuals, respectively (Figure 3.4-3). Based on these figures, numbers in MU1 in the Russian Arctic represent about 92% of the total flyway population and thus will be the major driver for changes in abundance at flyway level. Apart from the large

credibility intervals the estimates come with, this population has also continued to increase over the past years whereas the Baltic MU2 has been declining, and the North Sea MU3 has been fluctuating around 64,000 individuals since 2015. Some caution is needed for the estimates of the Baltic population, as data from Denmark is largely lacking. Surveys of breeding pairs in the Danish Baltic, however, do point at declines between 2018 and 2021, notably on the island of Saltholm between Denmark and Sweden (Nielsen et al. 2024).

Converted into breeding pairs, the posterior estimate for the size of the breeding population in the Russian MU1 in summer 2024 was 617,000 breeding pairs, thus exceeding 5.5 times the FRP and also exceeding the 200% threshold level by a large margin (Figure 3.4-4). The Baltic MU2 is well above (1.8 times) the FRP as well (an estimated 22,000 breeding pairs in 2024) but has now fallen below the 200% threshold. The North Sea MU3 was estimated at 19,000 breeding pairs, 1.6 times FRP (lower limit of 95% credibility interval just above FRP) but still below the 200% threshold of the FRP (Figure 3.4-4).




Figure 3.4-3. Top panel: January total flyway population counts (dots) of the Russia (br) population of Barnacle Goose, posterior means based on the IPM (solid line), 95% credible intervals (shaded area) and FRP as well as 200% of the FRP (dashed lines). Bottom panels: July population sizes of the three MU-populations along with posterior means and 95% credible intervals. Left (in red) MU1, centre (in blue) MU2, right (in green) MU3. Note the different scales on the y-axes. Further note that July counts of MU1 are not available and are estimated as latent variables within the IPM framework (and come with large 95% credibility intervals).




Figure 3.4-4. Posterior means (solid line) and 95% posterior intervals (shaded areas) for the number of breeding pairs in July for the three MU's of the Russia (br) population of Barnacle Goose, derived from the IPM. Dashed lines are the FRP as well as 200% of the FRP. Left (in red) MU1, centre (in blue) MU2, right (in green) MU3. In the IPM framework, the number of breeding pairs has been set as the number of individuals of 2 years and older, divided by 2. Note the different scales on the y-axes between MU1 and MU2/3.

b) Mortality and offtake

Survival rates derived from the IPM and combined for summer and winter, show that adults have much higher survival rates than juveniles (Figure 3.4-5, note that last year of the time series is based on incomplete data), which is also according to expectation. Adult survival in all three management units is estimated around 0.95, with little differences between the MUs (also given the credibility intervals). Especially in MU2 some variation seems to occur, for reasons unknown yet. Combined summer and winter survival for juvenile birds is estimated around 0.50 for the migratory birds in MU1 and MU2 while it is higher (0.58) for the sedentary birds in MU3. In all cases, the posterior credible intervals for juvenile survival are much wider than those for adult survival, and annual variation is more pronounced in all three MUs.

Combined offtake rates have increased over time for all populations (Figure 3.4-6; compare Figure 3.4-7 for national assessment). For MU1 offtake rates for adults and juveniles have stabilized around 3% and 6%, respectively. In MU2, there has been a continuous increase in both adult and juvenile offtake rates, now reaching a level around 10%. However, it should be noted that offtake in the Baltic population may be biased somewhat high, as numbers are divided among presumed MU1-birds and presumed MU2-birds. These cannot be distinguished for most of the year, and part of the derogations that are assigned to MU2 may actually affect MU1 (also being the dominant management unit regarding numbers). For MU3, offtake rates increased steeply after

2013, reflecting increase of management by the Dutch provinces in order to halt the ongoing increase of the breeding population and including round-up during wing moult in early summer. As a result, offtake rates for adults and juveniles even exceeded 30% in 2019/20 and 2020/21. But following the outcome of the EGMP-assessment in 2022, showing that numbers in MU3 were approaching FRP, derogation efforts in the Netherlands were reduced and numbers killed went down following the guidance given by the IWG. This is also clearly reflected in the figures presented in Figure 3.4-7, which show a considerable reduction in offtake under derogation from 2022 onwards. This reduction is solely the impact of reduced management during the summer period. During wintertime, derogations in the Netherlands are locally restricted due to provincial management policies that leave migratory birds partly untouched (e.g. by providing only restricted derogations or provide accommodation areas in which scaring and shooting is not allowed).

In 2025, data assessed for this report show that in the EU countries nearly 51,000 Barnacle Geese were killed under derogation, which is a slight increase from the 49,000 killed in 2023 (Figure 3.4-7), but still much lower than the numbers around 2020, when management effort to target summer population in the Netherlands reached its peak. In 2022-2024, 79% of all birds killed under derogation were killed in Denmark and the Netherlands. During the period 2022-2024, the use of derogations in Denmark has been slightly reduced whereas in Germany and Sweden there has been an increase. Earlier, an increase was also apparent in Finland, but national court cases nearly brought the number of derogation permits to zero in 2024.

Besides direct offtake, some countries also apply clutch management. This is mainly done in Germany (Schleswig-Holstein), where, in spring 2024, a derogation license to apply clutch management to up to 760 nests was issued, and actually 2322 eggs were treated to prevent hatching (data source European Environment Agency, see Figure 3.4-7). When assuming an average clutch size of 4 eggs, this would represent about 580 nests.




Figure 3.4-5. Posterior means and 95% confidence intervals for combined, i.e., summer and winter, juvenile (upper panel) and adult (lower panel) natural survival for geese in the three MUs of the Russia (br) population of Barnacle Goose. Left (in red) MU1, centre (in blue) MU2, right (in green) MU3. Note that this includes unknown offtake for the Russian population in Russia.




Figure 3.4-6. Posterior means (solid lines) and 95% posterior intervals (shaded area) for the combined derogation offtake rates of juveniles (top panels) and adults (bottom panels) for the three MU's of the Russia (br) population of Barnacle Goose. Left (in red) MU1, centre (in blue) MU2, right (in green) MU3.




Figure 3.4-7. Number of Barnacle Geese killed under derogation in EU countries during 2010-2023. Data retrieved from national agencies (the Netherlands, Germany, Denmark) or the data viewer of the European Environment Agency (<https://www.eea.europa.eu/en/analysis/maps-and-charts/derogations-and-exceptions-table-dashboards>). Only offtake by shooting or trapping is included (not clutch-management, see text).

c) Reproduction

The percentage of juveniles, reflecting a proxy for productivity in MU1 and MU2 (according to abundance it will mainly reflect productivity in MU1), in autumn flocks in (mainly) the Netherlands shows a high degree of annual variation. In most years, IPM-estimates are consistently higher, even if the recommended timing of age-ratio assessments in the field has been advanced since 2022 (to October-November), to avoid problems to identify juveniles (due to progress of plumage moult). Estimates for autumn 2024, both IPM-estimates and field data indicate that productivity was actually at a very low level. In the long-term data series from the Netherlands, it could be shown that productivity in the breeding season 2024 in the Russian Arctic was among the lowest since 1974/75. This was not only observed in Barnacle Geese but also other goose and swan species breeding in the same region as has been attributed to an exceptional cold spring and summer (Sovon 2025).

For MU2, productivity estimates based on the IPM show large fluctuations with some tendency to decline over time. Again, IPM estimates are higher than crude field data (of which some are even outside the 95% credibility intervals of the IPM estimates), but it should be noted that this data source is perhaps biased as it is mainly derived from assessment in the Helsinki metropolitan region in Finland, which is likely neither representative of other parts of Finland nor of other breeding populations in the Baltic, notably Sweden and Denmark (M. Mikkola-Roos & A. Lehtikoinen, pers. communication). Identification issues should not play a role in the field assessments, as these are done directly after the breeding season, when plumage characteristics of juveniles allow for a proper assessment.

Estimates for MU3 are higher than MU1 and MU2 and also show some increase over time (still dominated by large annual variation). Also in this MU, some of the field data is outside the 95% credibility interval of the IPM estimates and as in MU2 identification problems can be ruled out. However, it is not clear to what extent the sample taken in summer (both in the Netherlands and in North Rhine-Westphalia in Germany) is representative for entire MU3 (e.g. no field data are available in Schleswig-Holstein, where clutch management is in place).




Figure 3.4-8. Top panel: Observed autumn percentage of juveniles (dots) in the Russia (br) population of Barnacle Goose, posterior means (solid line) and 95% posterior intervals (shaded area). Bottom panels: Observed summer percentage of juveniles in the three MUs, along with posterior means and 95% posterior intervals. Left (red) MU1, centre (blue) MU2, right (green) MU3. Note that in MU1 there are no field data to compare with the IPM estimates (they are included in the assessments in autumn, given in the top panel).

3.4.6 Management guidance

The overall results of this year's assessment are broadly similar to those for 2024. The MU3 in Belgium, the Netherlands and Germany should still be subject to a coordinated derogation approach, in line with the 200% threshold set in the AFMP. The latest model output points at a population level which is again well above the size of the FRP (Figure 3.4-4), but below the 200% threshold. At present derogations affecting MU3 are only granted in the Netherlands. Derogations in Schleswig-Holstein in Germany are likely affecting MU2 and (even more so) MU1 only, given the time of the year in which they are undertaken. It remains unknown to what extent large-scale clutch management in Schleswig-Holstein affects total numbers in July and productivity in MU3. In Niedersachsen, no derogation takes place, and Belgium considers its small breeding population to be of feral

origin (Nagy et al. 2021b, F. Verhagen pers. communication), so is left out of this coordination. In the Netherlands, a coordinated approach among the provinces (which are each responsible for their own goose management) has been installed in order to facilitate implementation of the AFMP and avoid numbers falling below the FRP. To facilitate this process, the FRPs for the Netherlands in the AFMP have been divided over the 12 provinces (Sovon 2022) and management is assessed annually by the 12 regional wildlife councils.

Worth noticing is the fact that the current IPM-estimates indicate that MU2 has now also fallen below the 200% threshold level and would thus require coordination among Range States (notably Denmark, Sweden, and Finland), as far as actions affecting local breeding populations are undertaken or planned. Based on derogation data collected so far, offtake in these countries likely affect mainly MU1, but there is still a need to investigate a more data-based division of offtake under derogation in these countries (perhaps differentiating among regions or time of the year), to evaluate to what extent birds from MU2 might be affected.

Regarding MU1, the results indicate that this population has increased until very recently and that its current population level is way beyond the 200% threshold. Breeding opportunities in the Russian Arctic are also assumed to expand, as shown by Lameris et al. (2023) for Novaya Zemlya, as a result of climate change and increase in available breeding habitat. Whether the similar numbers for summer 2023 and 2024 (Figure 3.4-3 and 3.4-4) are signs of a stabilisation is too early to assess. Low productivity in the Russian Arctic in 2024 has been attributed to adverse weather conditions during the breeding season (see above). It should be noted though, that the overall flyway population, as assessed in January, and largely dominated by the Russian breeding population, currently does not show an eminent growth.

There is no indication that highly pathogenic avian influenza (HPAI) has resulted in considerable declines, as was initially observed in the Svalbard population of Barnacle Goose. It is unknown if HPAI, among other factors, may have slowed down the earlier population increase. Caliendo et al. (2024) estimated that in 2020/21 and 2021/22 up to 4.8% and 7.4% of the Barnacle Goose present in the Netherlands died of avian influenza. During 2024/25, at least in the Netherlands, higher mortality was again observed among wintering birds but apparently involving fewer birds than during 2020-2022. In the most recent winter, 2025/26, HPAI-casualties of Barnacle Geese were still found, but again numbers seemed to be smaller than in the initial winters 2020-2022, and other species (e.g. Greylag Goose) seemed to be more severely affected.

In terms of monitoring data for the IPM, we can currently derive good winter census results from the extensive IWC and goose counting networks in the range states. It would be desirable to make data for Germany (notably Schleswig-Holstein) available earlier, so all German data could feed into the IPM-calculations in time. Also, regarding productivity, a large sample is collected in autumn in the Netherlands, but preferably this should be extended to Germany, Denmark and Sweden to sample the newly established wintering areas (which potentially may consist of different birds and different rates of productivity).

Regarding data collected in the summer period there are larger and more profound gaps, especially in MU2. At present, Finland is the only country in which comprehensive (late) summer counts have been established since 2008. For Sweden, where summer census data were lacking for nearly the entire period of interest, we have now used results from the mid-September count instead. This count is carried out before migratory birds from MU1 arrive, and it is only two weeks after the summer count in Finland (which takes place by the end of August). It is assumed that in this short period, transition rates between the two countries are low, but this assumption should preferably be underpinned by data from resightings of ringed birds or tracking data (which is in progress as part of a Swedish project on Barnacle Geese). Data from resightings of ringed birds do confirm that in both Sweden and Finland the summer counts are carried out before arrival of Russian birds from MU1 (L. de Vries, pers. communication). Besides, given the large fluctuations in the count results for September in Sweden, it is recommended to check whether the coverage of the counts does affect the final results, or if other explanations could be found for some years with low numbers (e.g. earlier emigration to Denmark or other countries further south). This is especially important, as the current data suggests that the numbers in MU2 are below 200% of the FRP and the modus operandi in the AFMP calls for a coordinated derogation effort (concerning offtake that may

affect local breeding populations). Relevant to note in this context is also that breeding numbers in Sweden have declined in the period 2007-2018³. So far, data has only periodically been collected in the Oslofjord area in Norway and in Denmark and it is recommended to continue these periodical counts, especially in Denmark, where breeding bird surveys suggest a decline in the number of breeding pairs, at least at local level (Nielsen et al. 2024). Counts have been carried out in Denmark in 2025 but have not yet been processed due to financial constraints. In MU3, it would be highly desirable to include up-to-date counts from Schleswig-Holstein (now available with considerable delay), along with data on productivity (age-ratio counts in summer) and a more timely assessment and publication of these data.

Productivity data from MU2 heavily depend on the Helsinki area in Finland, but additional data have been collected in other regions and will be added in the assessment in 2027. Still, productivity data from especially Sweden and Denmark would allow a more comprehensive assessment of breeding success in the entire Baltic region.

Furthermore, seasonal and MU assignment of offtake still depends on some assumptions and expert judgement, as most data are only available as a total figure for the entire calendar year and assignment to MU-level is challenging. Currently, the Netherlands is the only country with a monthly data resolution, allowing us to make more precise assignment of derogation figures to each respective MU (notably segregating migratory and sedentary populations to a large extent). Also, data from Denmark partly allows this. However, derogations granted often do not take into account seasonal occurrence of birds from different MUs and may even refer to different calendar years (e.g. from July to June). This aspect needs further attention in the preparations for the next IPM-assessments. If Range States within MU2 would investigate their derogation data in detail to discern to what extent birds from MU2 are affected (see above), this would also be beneficial to feed into the IPM in order to carry out a more precise assessment.

³ See https://nature-art12.eionet.europa.eu/article12/summary?period=3&subject=Branta+leucopsis&reported_name=.

3.5 East Greenland breeding population of Barnacle Goose *Branta leucopsis*

3.5.1 Range States and Management Units

The Range States for the *E. Greenland/Scotland & Ireland population of Barnacle Goose* include Greenland, Iceland, Republic of Ireland and United Kingdom (Figure 3.5-1). The population is managed as one Management Unit (MU) (Jensen et al. 2018, Nagy et al. 2021a).




Figure 3.5-1. Annual distribution and migration routes for the *E. Greenland/Scotland & Ireland population of Barnacle Geese*, including breeding (red), staging (green) and wintering (blue) areas.

3.5.2 Population FRPs and targets

The FRP for the breeding season is 19,400 pairs (Nagy et al. 2021a). The FRP for the entire population has been set at 54,000 wintering individuals. Being an Annex 1 species of the EU Birds Directive, the AFMP does not aim to maintain the population at a certain target level. In EU countries (Ireland) and the UK management is carried out under the conditions for derogation, outlined in Art. 9 of the EU Birds Directive. Furthermore, the species is strictly protected under the Bern Convention. There are open hunting seasons for the species in Iceland (which has entered a reservation in respect of the Bern Convention's Appendix II listing of Barnacle Geese) and Greenland.

3.5.3 Management strategies

The AFMP aims to prevent the population declining below the defined FRPs (Nagy et al. 2021a). Thus, the FRPs represent the lower limit of the legally acceptable population size but does not reflect targets for population reduction. Monitoring of the population size and harvest, and predictive modelling of the cumulative impact of national derogation measures and hunting are used to inform national decision-making to ensure the population remains above the FRPs. The cumulative impact of derogation and hunting and the non-lethal measures taken to

prevent damage/risk on the population are assessed periodically, along with the likelihood of serious damage to agriculture and risk to air safety and to other flora and fauna (including the Arctic ecosystems), as well as the effectiveness of these.

Within this framework, it has also been agreed to coordinate monitoring of the population and offtake under derogations and hunting when the actual size of the populations is below 200% of the defined FRP. This includes prediction of population development, coordination of offtake and taking coordinated conservation measures, where necessary. Note, however, that the population size has perhaps never exceeded 200% of the FRP.

As the population is apparently approaching the FRP, it was agreed at IWG9 in June 2024 that Iceland and the United Kingdom should seek agreement on the maximum level of offtake to be permitted (if any) and the split between the two Range States, and further develop and implement a coordination mechanism to ensure adherence to these limits. Since 2024, Iceland and the United Kingdom have informed the EGM IWG in writing on the agreed levels of offtake, the agreed coordination mechanism, and reported on the implementation and adherence to the agreed levels of offtake.

3.5.4 Assessment protocol

In 2020, NatureScot and the Department of Housing, Local Government and Heritage Ireland, funded the development of an integrated population model (IPM) for the purpose of better understanding the population dynamics of the flyway population of E Greenland (br) barnacle geese and in order to inform the management of offtake for the species.

We refer to McIntosh et al. (2023) for the following description of the IPM, which is a pre-breeding census model with an annual time-step and anniversary date in March. Annual change in March abundance is described as:

$$N_{t+1} = N_t \theta \left\{ \begin{array}{l} (1 - h_t^i) \left((1 - p_t^{Islay}) + p_t^{Islay} (1 - h_t^s) \right) + \\ r_t (1 - v^i h_t^i) \left((1 - p_t^{Islay}) + p_t^{Islay} (1 - v^s h_t^s) \right) \end{array} \right\}$$

where N_t is the March population size at time t , θ is the constant rate of natural survival, p_t^{Islay} is the proportion of the March flyway population on Islay at time t , h_t^i is the annual harvest rate in Iceland, h_t^s is the annual harvest rate in Scotland, v^i is the differential vulnerability of juveniles in Iceland, v^s is the differential vulnerability in Scotland, and r_t is the pre-season age ratio (juvenile: adult ratio at the start of the hunting season).

To model annual change in March abundance we assumed that: a) harvest occurs sequentially (first in Iceland, then in Scotland), b) differential vulnerability of juveniles in Scotland is constant throughout the winter (Calvert et al. 2017), c) natural mortality is distributed evenly throughout the year (Gauthier et al. 2001). Lastly, we assumed that shooting mortality is additive to natural mortality as observed in numerous other goose populations (Gauthier et al. 2001, Sedinger et al. 2007, Cooch et al. 2014, Koons et al. 2014).

We assume six months of natural mortality to predict pre-hunting population size:

$$N_t^F = N_{A,t}^S \theta^{6/12} + N_{A,t}^S \theta^{6/12} r_t$$

where N_t^F is the autumn population size and $N_{A,t}^S$ is the adult spring population size.

Harvest occurs first in Iceland (H^i) in the early autumn:

$$H_t^i = N_{A,t}^S \theta^{6/12} h_t^i (1 + r_t v^i)$$

To estimate Scottish harvest (H^s) we assume an additional month of natural mortality and that individuals survive harvest in Iceland. Winter derogation shooting occurs predominantly on Islay, therefore only Islay-wintering birds experience Scottish shooting mortality.

Number surviving Iceland harvest is:

$$(N_t^F - H_t^I)\theta^{1/12} = N_{A,t}^S\theta^{7/12}(1 - h_t^i) + N_{A,t}^S\theta^{7/12}r_t(1 - v^i h_t^i)$$

Scottish harvest (H^S) is then

$$H_t^S = N_{A,t}^S\theta^{7/12} \left\{ \left(p_t^{Islay} \left((1 - h_t^i) h_t^s \right) \right) + r_t \left(p_t^{Islay} \left((1 - v^i h_t^i) v^s h_t^s \right) \right) \right\}$$

We estimated annual harvest rates for different age classes.

Adults (h_t^A):

$$h_t^A = h_t^i + \theta^{1/12} \left(p_t^{Islay} \left((1 - h_t^i) h_t^s \right) \right)$$

Juveniles (h_t^J):

$$h_t^J = v^i h_t^i + \theta^{1/12} \left(p_t^{Islay} \left((1 - v^i h_t^i) v^s h_t^s \right) \right)$$

Annual survival rate (s_t) is derived from apparent natural survival (q) and harvest mortality (h_t). Due to an absence of data on unretrieved harvest, crippling losses (unobserved harvest mortality) are implicitly included in the estimate of natural mortality. Adult survival rate is:

$$s_t^A = \theta(1 - h_t^A)$$

and juvenile survival rate is:

$$s_t^J = \theta(1 - h_t^J)$$

Raw data and the results of the 2026 update of the IPM are available from the [EGMP Data Centre](#).

3.5.5 Status

a) Abundance

After a peak population of 80,000 in 2006 and in 2012, posterior estimates of flyway abundance declined to 65,516 (55,558 – 76,459) in March 2026 (Figure 3.5-2). For much of the period of record, abundance on Islay exceeded that in all other wintering areas, but that pattern has been reversed since 2018.




Figure 3.5-2. Development of the March population size of E. Greenland/Scotland & Ireland Barnacle Geese as based on the IPM. The FRP = 54 thousand. Shading represents the 95% credible intervals.

b) Mortality and offtake

Natural survival (i.e., 1 – the natural mortality rate) was relatively high and stable until 2007 when it became more variable, with unusually low natural survival during 2007 – 2009, in 2017, and during 2020 – 2021 (Figure 3.5-3). The latter period of low survival might be attributed to an outbreak of avian influenza, but it is difficult to say whether survival was in fact lower than is typical because of the wide credible intervals.




Figure 3.5-3. Natural survival rates (i.e., 1 – the natural mortality rate) of E. Greenland/Scotland & Ireland Barnacle Geese as based on the IPM. Shading represents the 95% credible intervals.

The total harvest rate of adults has increased over the period of record, from around 0.01 to a peak of 0.05 (0.03 – 0.07) in 2017 (Figure 3.5-4). Thereafter, harvest rate declined to 0.02 (0.01 – 0.03) in 2025. Annual survival rate of adults (including both harvest and natural mortality) declined at the same time harvest rates were increasing, suggesting that harvest may have contributed to the decline in flyway abundance (although other factors cannot be ruled out).




Figure 3.5-4. Adult harvest rates (left) and annual survival rates (right) of E. Greenland/Scotland & Ireland Barnacle Geese as based on the IPM. Shading represents the 95% credible intervals.

Recreational harvest (and overall harvest rate) in Iceland has generally increased over the period of record (Figure 3.5-5, Figure 3.5-6) but substantially decreased in 2025 with an estimated harvest of 1,681 (1,017 – 2,589) individuals. In Scotland, derogations increased starting in 2012 in response to a plan to limit agricultural conflicts but has now been reduced to near zero in response to avian influenza and the observed population decline. The 2025 estimated harvest in Scotland was 233 (54 – 628) individuals.




Figure 3.5-5. Offtake of E. Greenland/Scotland & Ireland Barnacle Geese as based on the IPM. Shading represents the 95% credible intervals.




Figure 3.5-6. Harvest rates (including derogations) of E. Greenland/Scotland & Ireland Barnacle Geese as based on the IPM. Shading represents the 95% credible intervals.

c) Reproduction

The post-breeding age ratio has been moderately variable over time, although perhaps somewhat lower since 2006 than previously (Figure 3.5-7). It is possible that this variability in reproduction, along with the lower annual survival rates in more recent years, could have contributed to the more recent decline in flyway population size (Figure 3.5-4).




Figure 3.5-7. Post-breeding age ratio of E. Greenland/Scotland & Ireland Barnacle Geese as based on the IPM. Shading represents the 95% credible intervals.

3.5.6 Management guidance

It appears that a combination of factors, possibly including decreased productivity, as well as increased harvest and natural mortality rates, may have been responsible for the decline in flyway abundance over the last decade. Given the harvest rates in 2025, estimates from the IPM give a 1% probability that the March 2026 population is below the FRP of 54,000, based on the proportion of 2026 abundance estimates falling below this value through the total number (60,000) of model iterations. However, because of the proximity of the population to the FRP over the past several years, the Adaptive Flyway Management Plan requires tighter coordination of offtake between Iceland and Scotland to ensure the population does not fall below the FRP.

Table 3.5-1 provides a range of scenarios of varying levels of offtake applied to varying spring population sizes to determine the probability (based on the IPM) of the population falling below the FRP the following year. Thus, this table expresses the risk of falling below the FRP for a range of population sizes and levels of offtake *for any given year*. Once an acceptable risk level is established, the table can provide the **maximum** acceptable offtake for any population size. Probabilities account for uncertainty in natural mortality and reproductive rates, as well as for sampling error in estimated population size. Probabilities are updated each time the IPM is updated.

Table. 3.5-1. Approximate probability that the following year's population size of Greenland Barnacle Geese is lower than the FRP of 54k for varying levels of population size and offtake in any given year. The table does not refer to any particular year. March population sizes (in thousands [k]) for any focal year (t) are provided in the leftmost column and varying offtake levels (in thousands [k]) are represented in the top row. Values in each coloured cell represent the probability that the population size in the following year (i.e., t + 1 will be below the FRP, $P(N_{t+1} < 54k)$). Colour scale represents increasing level of risk, where warmer colours indicate higher probabilities that the population will fall below the FRP.

March pop (k)	Offtake (k)																								
	0	0.25	0.5	0.75	1	1.25	1.5	1.75	2	2.25	2.5	2.75	3	3.25	3.5	3.75	4	4.25	4.5	4.75	5	5.25	5.5	5.75	6
54	0.29	0.31	0.32	0.34	0.35	0.36	0.38	0.39	0.41	0.43	0.44	0.46	0.47	0.49	0.51	0.52	0.55	0.56	0.57	0.59	0.60	0.62	0.63	0.65	0.66
55	0.23	0.25	0.26	0.27	0.29	0.30	0.32	0.33	0.35	0.37	0.38	0.39	0.40	0.43	0.44	0.46	0.47	0.49	0.51	0.52	0.54	0.55	0.57	0.58	0.59
56	0.18	0.20	0.21	0.23	0.23	0.24	0.25	0.28	0.28	0.31	0.32	0.34	0.35	0.37	0.38	0.39	0.41	0.42	0.44	0.46	0.47	0.48	0.50	0.51	0.53
57	0.14	0.15	0.17	0.17	0.19	0.20	0.21	0.22	0.23	0.24	0.26	0.27	0.29	0.30	0.32	0.33	0.35	0.36	0.37	0.39	0.40	0.42	0.44	0.45	0.46
58	0.11	0.12	0.13	0.14	0.14	0.16	0.16	0.18	0.19	0.19	0.21	0.22	0.24	0.24	0.26	0.27	0.29	0.30	0.32	0.32	0.34	0.36	0.37	0.39	0.40
59	0.08	0.09	0.10	0.10	0.11	0.12	0.13	0.14	0.15	0.16	0.17	0.18	0.19	0.20	0.21	0.22	0.24	0.25	0.26	0.28	0.29	0.31	0.31	0.32	0.34
60	0.06	0.07	0.08	0.08	0.09	0.09	0.11	0.11	0.12	0.12	0.13	0.14	0.15	0.16	0.17	0.18	0.19	0.20	0.22	0.22	0.23	0.25	0.27	0.28	0.29
61	0.05	0.05	0.05	0.06	0.06	0.07	0.08	0.08	0.09	0.09	0.10	0.11	0.12	0.12	0.13	0.14	0.15	0.15	0.18	0.18	0.19	0.20	0.21	0.23	0.25
62	0.03	0.03	0.04	0.04	0.05	0.05	0.06	0.06	0.06	0.07	0.08	0.08	0.09	0.10	0.10	0.11	0.12	0.13	0.13	0.14	0.16	0.16	0.17	0.19	0.19
63	0.02	0.02	0.03	0.03	0.03	0.04	0.04	0.04	0.05	0.05	0.06	0.06	0.07	0.08	0.08	0.08	0.10	0.10	0.10	0.11	0.12	0.13	0.14	0.15	0.15
64	0.02	0.02	0.02	0.02	0.03	0.03	0.03	0.03	0.04	0.04	0.04	0.04	0.05	0.06	0.06	0.07	0.07	0.08	0.08	0.09	0.09	0.10	0.10	0.12	0.12
65	0.01	0.01	0.01	0.01	0.02	0.02	0.02	0.03	0.03	0.03	0.03	0.03	0.04	0.04	0.04	0.05	0.05	0.06	0.06	0.07	0.07	0.08	0.08	0.09	0.10
66	0.01	0.01	0.01	0.01	0.01	0.01	0.02	0.02	0.02	0.02	0.02	0.03	0.03	0.03	0.04	0.04	0.04	0.04	0.05	0.06	0.06	0.06	0.07	0.08	0.08
67	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.02	0.02	0.02	0.02	0.03	0.03	0.03	0.04	0.04	0.04	0.04	0.05	0.05	0.06
68	0.00	0.00	0.00	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.02	0.02	0.02	0.02	0.02	0.03	0.03	0.03	0.04	0.04	0.04
69	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.02	0.02	0.02	0.02	0.03	0.03	0.03
70	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.02	0.02	0.02	0.02	0.02	0.02	0.02

Appendix A – Data overview

A.1. Pink-footed Goose – Svalbard (br) population

[This data be made available on 8 June 2026]

A.2. Taiga Bean Goose

[This data be made available on 1 June 2026]

A.3. Greylag Goose – NW Europe (br) population

Table A.3.1 Overview of available monitoring data in the NW Europe (br) Greylag Goose population.

Grey cells mark data for MU1 and blue cells for MU2.

X = data collected annually/regularly and reported to EGMP, x = data collected annually/regularly, (x) = data collected in part of the country and/or not annually, - = no data collected or reported to the EGMP, * = 0 or not relevant range state in this respect.

	NO	SE	FI	DK	DE	NL	BE	FR	ES
Population counts in January (received through IWC or derived from national goose counts)	X	X	*	X	(x) ¹	X	X	X	(x) ²
Summer count	(x) ³	X ⁴	X ⁵	X ⁶	(x) ⁷	X	X	(x) ⁸	*
Productivity	(x) ⁹	(x) ⁹	(x) ⁹	-	(x) ⁹	(x) ⁹	-	-	-
Hunting bag	X	X	X	X	X ¹⁰	*	X	X ¹¹	X
Split hunting data into March-Aug and Sep-Feb	(x) ¹²	(x) ¹³	(x) ¹⁴	(x) ¹⁵	-	*	(x) ¹⁶	X	X
Derogation	(x) ¹⁷	X	X	X	(x) ¹⁸	X	X	*	*
Split derogation data into March-Aug and Sep-Feb	(x) ¹⁹	(x) ²⁰	(x) ²¹	X	-	X	(x) ²¹	*	*
Crippling rate		(x) ²²				(x) ²²			

- 1) Available from IWC most years, but the coverage is unknown.
- 2) Available from IWC most years, but the coverage is limited.
- 3) Country-wide estimate from 2022 has been made available. Future count/estimate interval unknown.
- 4) September count is used. Coverage could be improved, and counts do not account for hunting and migration.
- 5) To estimate population size, organized counts have been carried out in 2022 and 2023. GPS-tracking has been used to distinguish between birds from the C and NW Europe (br) populations.
- 6) Counted biennially in August.
- 7) Available from Nordrhein-Westfalen (since 2011) and Niedersachsen (2018-2023). Data from Schleswig-Holstein is available for June and September 2018-2023.
- 8) Available every 6 years from 2022.
- 9) Samples of age ratios from selected sites in N and S Norway provided. See Figure 3.3-10 and the [EGMP Database](#) for details.
- 10) Data Source: Datenspeicher Jagd Eberswalde, Thünen-Institut.
- 11) Method unknown.
- 12) Hunting season opens from 21.07. Assume all hunting takes place Sep-Feb.
- 13) Open hunting season 11.08-31.01. Assume all hunting takes place Sep-Feb. Conditional hunting season: all year, but assume all takes place between March-Aug.
- 14) Hunting season 10.08-31.12. Assume all hunting takes place Sep-Feb.
- 15) Hunting season 01.08-31.01. Assume all hunting takes place Sep-Feb.

- 16) Hunting season 15.07-31.01, but 15.07-14.08 and 01.10-31.01 constrained to prevent (crop) damage in the absence of other satisfying solutions. Open hunting season 15.08-30.09. Assume all hunting takes place Sep-Feb.
- 17) No routine data collection, but few individuals (~1000).
- 18) Available in most years.
- 19) All year, assume all derogation takes place between March-Aug.
- 20) Derogation period: 01.01-09.08, the majority takes place in July-Aug. Assume all derogation takes place between March-Aug.
- 21) Assume all derogation takes place between March-Aug.
- 22) Not collected annually, and only for part of the flyway.

Table A.3.2 Overview of model parameters and their source

Parameter	Description	Value	Source
Φ	annual survival in absence of hunting	0.88	allometric relationship (Johnson et al. 2012)
$\phi(0.90)$	annual survival of young from MU1	0.79	loosely based on Pistorius et al. (2006) and Schneider & Bacon (2022)
α	annual rate of production of young by birds aged 3+	0.46	derived using ϕ and population growth rate of 1.014 from EGMP Population Status and Offtake Assessment Report (2022)
ψ_1	proportion of MU1 birds wintering in the North	0.67	based on marking data (Leo Bacon, pers. comm.)
ψ_2	proportion of MU2 birds wintering in the North	0.95	based on marking data (Leo Bacon, pers. comm.)
π_1, π_2	fidelity of MU1 and MU2 birds	1.0	Schneider & Bacon (2022), recognizing that lack of fidelity is typically temporary
$\begin{bmatrix} n_{1,1} \\ n_{2,1} \\ n_{3,1} \\ n_{1,2} \\ n_{2,2} \\ n_{3,2} \end{bmatrix}$	initial population sizes (in thousands) in fall 2022, where the first subscript denotes age and the second denotes MU	$\begin{bmatrix} 72.2 \\ 56.4 \\ 201.1 \\ 120.3 \\ 94.0 \\ 334.9 \end{bmatrix}$	derived based on estimates of breeding pairs in 2018 (Szabolcs Nagy, pers. comm.) and the stable age distribution of the matrix model in the absence of harvest
h_{ijk}	rate of offtake of age i , season j , and area k	0.0 to 0.4 in increments of 0.02	simulated to project population sizes in 2030
ν_s	differential vulnerability of young in summer	1.0	assumed given no selectivity in summer derogations
ν_w	differential vulnerability of young in winter	2.0	assumed to be similar to pink-footed geese (Johnson et al. 2020)

A.4. Barnacle Goose – Russia breeding population

Table A.4. Overview of available monitoring data in the Russia (br) Barnacle Goose population.
X data collected at national level/annually, *(x)* data collected but not annually and/or not at national level, - data currently not collected, * not relevant range state in this respect.

	RU	FI	EE	SE	NO	DK	DE	NL	BE	Remark
January census	*	*	*	X	*	X	X ¹	X	X	Timely publication of data collected in Schleswig-Holstein (DE) would facilitate assessment of winter counts.
Summer census	-	X	-	X ²	(x)	(x)	(x)	X	X	Interaction between counts in SE and FI (and DK) needs to be investigated. Timely processing of count data from Denmark would facilitate the annual update of the IPM (most recent data from 2025 was not available).
Productivity, MU1 and MU2	*	*	*	-	*	-	X	X	-	Autumn, Oct-Dec. Data from DE, DK, and SE would be valuable.
Productivity, MU2	*	(x)	-	-	(x)	-	*	*	*	Summer, Jul-Aug. Inclusion of more comprehensive productivity data in FI in 2027 is foreseen. Recommendation to extent productivity assessments to SE and DK.
Productivity, MU3	*	*	*	*	*	*	(x)	X	-	Summer, Jul. DE only North-Rhine Westphalia. Preferably inclusion of data from other parts of DE (Niedersachsen, Schleswig-Holstein)
Offtake, hunting	-	*	*	*	*	*	*	*	*	In EU-countries only derogations.
Offtake, derogations	*	X	X	X	X ³	X	X	X	X	Mostly annual totals (apart from monthly data in NL), more detailed data requested especially from SE in order to improve estimates for individual MUs.

¹ Note that Germany only submits data based on published count results.
² For Sweden, the national count in mid-September is used as a proxy for numbers in summer.
³ Norway is not an EU Member State but applies similar rules when it comes to management of Barnacle Goose, although derogations are for scaring purposes only.

A.5. Barnacle Goose – East Greenland breeding population

Table A.5. Overview of available monitoring data in the E Greenland (br) Barnacle Goose population.
X data collected (nearly) annually and reported to EGMP, *x* data collected (nearly) annually, *(x)* data collected in part of the country and/or not annually, - no data collected or reported to the EGMP, * 0 or not relevant range state in this respect.

	UK	Ireland	Iceland	Greenland
Flyway total every 3 years	X	X	*	*
Islay March count - annual	X	*	*	*
Other totals in Scotland - annual	X	*	*	*
Breeding bird count in Iceland every 3 year	*	*	X	*
Offtake	X	X	X	- / (x)
Productivity	X	-	-	-
Wings	X	*	X	*
Survival	-	-	-	-

References

- Alhainen M., Kanstrup N., Piironen A., Heinicke T., Hall, K., Sørensen I.H., Madsen J., Germain R. and Lewis, M. (compilers) 2025. International Single Species Action Plan for the Conservation of the Taiga Bean Goose *Anser fabalis fabalis* (Scandinavia (br), Finland & NW Russia (br), Germany & Poland (nbr), C Asia (nbr) populations). AEWA Technical Series No. 80. Bonn, Germany.
- Aubry, P., M. Guillemain, G. H. Jensen, M. Sorrenti, and D. Scallan. 2020. Moving from intentions to actions for collecting hunting bag statistics at the European scale: some methodological insights. *European Journal of Wildlife Research*, 66,70 (2020), <https://doi.org/10.1007/s10344-020-01400-2>
- Bairlein, F., J. Dierschke, V. Dierschke, V. Salewski, O. Geiter, K. Hüppop, U. Köppen, and W. Fiedler. 2014. Atlas des Vogelzuges. Ringfunde deutscher Brut- und Gastvögel. AULA-Verlag, Wiebelsheim.
- Baveco, H., P. W. Goedhart, K. Koffijberg, H. van der Jeugd, L. de Vries, and R. Buij. 2021. Development of an integretaed population model for Barnacle Geese of the Russian management unit'. Progress Report prepared by the Dutch working group on Barnacle Goose in collaboration with the EGMP Data Centre. in S. Nagy, H. Heldbjerg, G. H. Jensen, F. A. Johnson, J. Madsen, O. Therkildsen, E. Meyers and S. Dereliev (eds.), Netherlands Population of the Barnacle Goose (*Branta leucopsis*). AEWA EGMP Programme No. 3. Bonn, Germany.
- Blüml, V. & H. Kruckenberg 2023. Gänsemonitoring und Gänsemanagement in Niedersachsen. Naturschutz und Landschaftspflege 51, NLWKN, Hannover.
- Buij, R., Moonen, S., Müskens, G. et al. (2026). Crippling rates of waterfowl by gunshot in Northwestern Europe: interspecific and geographic differences give clues to possible management challenges and options. *European Journal of Wildlife Research* 72, 35. <https://doi.org/10.1007/s10344-026-02070-2>
- Caliendo, V., E. Kleyheeg, N. Beerens, C.J. Camphuysen, K., R. Cazemier, A.R.W. Elbers, R.A.M Foucher, L. Kelder, T. Kuiken, M. Leopold, R. Slaterus, M.A.H. Spierenburg, H. van der Jeugd, H. Verdaat, and J.M. Rijks. 2024. Effect of 2020–21 and 2021–22 Highly Pathogenic Avian Influenza H5 Epidemics on Wild Birds, the Netherlands. *Emerging Infectious Diseases*, 30(1), 50-57. <https://doi.org/10.3201/eid3001.230970>.
- Calvert, A. M., Alisaukas, R. T., & White, G. C. 2017. Annual survival and seasonal hunting mortality of midcontinent snow geese. *The Journal of Wildlife Management*, 81(6), 1009–1020.
- Clausen, K.K., T.E. Holm, L. Haugaard and J. Madsen. 2017. Crippling ratio: A novel approach to assess hunting-induced wounding of wild animals. *Ecological Indicators*, 80: 242-46, <https://doi.org/10.1016/j.ecolind.2017.05.044>
- Cooch, E. G., Guillemain, M., Boomer, G. S., Lebreton, J., & Nichols, J. D. 2014. The effects of harvest on waterfowl populations. *Wild*, 4, 220–276.
- EU. 2020. Composite European Commission Report on Derogations according to article 9 of Directive 2009/147/EC on the Conservation of Wild Birds.
- Fox, A.D. & J.O. Leafloor. (eds.) 2018. A global audit of the status and trends of Arctic and Northern Hemisphere goose populations. Conservation of Arctic Flora and Fauna International Secretariat:

Gauthier, G., R. Pradel, S. Menu, J.-D. Lebreton. 2001. Seasonal survival of Greater Snow Geese and effect of hunting under dependence in sighting probability. *Ecology*, 82(11): 3105–3119, [https://doi.org/10.1890/0012-9658\(2001\)082\[3105:SSOGSG\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2001)082[3105:SSOGSG]2.0.CO;2)

Gupte, P., K. Koffijberg, G. Mskens, M. Wikelski, and A. Klzsch. 2019. Family size dynamics in wintering geese. *Journal of Ornithology*, 160: 363-75, <https://doi.org/10.1007/s10336-018-1613-5>

Haas, F., H. Kampe-Persson & L. Nilsson. 2023. Inventering av höstrastande och övervintrande gäss i Sverige – årsrapport för 2022. Biologiska institutionen, Lunds universitet.

Haas, F. 2025. Antalet höstrastande och övervintrande gäss i Sverige 2024/2025. 2025. Biologiska institutionen, Lunds universitet.

Heldbjerg, H., F. A. Johnson, K. Koffijberg, R. McKenzie, S. Nagy, G. H. Jensen and J. Madsen. 2021. Population Status and Assessment Report 2021. AEWA EGMP Technical Report No. 19. Bonn, Germany.

Hornman, M., M. Kavelaars, K. Koffijberg, F. Hustings, E. van Winden, P. van Els and R. Kleefstra. 2021. Watervogels in Nederland in 2018/2019. Sovon rapport 2021/01, RWS-rapport BM 21.08. Sovon Vogelonderzoek Nederland, Nijmegen.

Hornman M., Koffijberg K., van Oostveen C., van Winden E., Louwe Kooijmans J., Kleefstra R. and Soldaat L. 2024. Watervogels in Nederland in 2022/2023. Sovon-rapport 2024/96, RWS-rapport BM 24.39. Sovon Vogelonderzoek Nederland, Nijmegen. <https://stats.sovon.nl/pub/publicatie/22231>

Jensen, G.H., H. Heldbjerg, I.H. Sørensen, P. Clausen and C.L. Pedersen. 2023a. Summer count of Greylag Geese in Denmark 2022. Aarhus University, DCE - Danish Centre for Environment and Energy, 27 s. – Scientific briefing no. 2023|1

Jensen, G.H., F.A. Johnson, H. Baveco, K. Koffijberg, P.W. Goedhart, R. McKenzie and J. Madsen. 2022. Population Status and Assessment Report 2022. AEWA EGMP Technical Report No. 20 Bonn, Germany.

Jensen, G. H., J. Madsen, S. Nagy, and M. Lewis. 2018. AEWA International Single Species Management Plan for the Barnacle Goose (*Branta leucopsis*) - Russia/Germany & Netherlands population, East Greenland/Scotland & Ireland population, Svalbard/South-west Scotland population. AEWA Technical Series No. 70. Bonn.

Jiguet, F., Mayrose, A., Piha, M., Efrat, R., Seimola, T., Sapir, N., Timonen, S., Wasserlauf, Y., Besnard, A., Bocher, P., Kölzsch, A., Lehtikoinen, A., Monnet, A.-C., Procházka, P., and Perlman, Y. 2025. GNSS spoofing in conflict zones disrupts wildlife tracking and hampers research and conservation efforts. *Nature Communications* 16: 1199. <https://doi.org/10.1038/s41467-025-56630-2>

Johnson, F. A., H. Heldbjerg, S. Nagy, and J. Madsen. 2021. Setting population-size targets for geese causing socio-economic conflicts. *Ambio* 51:209–225, <https://doi.org/10.1007/s13280-021-01539-5>

Johnson, F.A., I.H. Sørensen, H. Baveco, K. Koffijberg, R.R. Germain, and J. Madsen. 2024. Population Status and Assessment Report 2024. EGMP Technical Report No. 22. Bonn, Germany.

Johnson, F.A., J. Madsen, K. K. Clausen, M. Frederiksen and G.H. Jensen. 2022. Assessing the value of monitoring to biological inference and expected management performance for a European goose population. *Journal of Applied Ecology* 60(1):132-145, <https://doi.org/10.1111/1365-2664.14313>

- Johnson, F. A., and K. Koffijberg. 2021. Biased monitoring data and an info-gap model for regulating the offtake of greylag geese in Europe. *Wildlife Biology*, 1:wlb.00803, <https://doi.org/10.2981/wlb.00803>
- Johnson, F. A., M. A. H. Walters, and G. S. Boomer. 2012. Allowable levels of take for the trade in Nearctic songbirds. *Ecological Applications* 22:1114–1130, <https://doi.org/10.1890/11-1164.1>
- Jones, J. 2026. Making sense of the 2025-26 Taiga Bean Goose influx. BirdGuides. <https://www.birdguides.com/articles/making-sense-of-the-2025-26-taiga-bean-geese-influx/>.
- Koffijberg, K. 2022. Bruterfolgsmonitoring bei Gänsen und Schwänen: Bestimmung von Alt- und Jungvögeln. Version 15.10.2022. Dachverband Deutscher Avifaunisten, <https://austausch.dda-web.de/s/Bestimmung-GuS-Jungvoegel>.
- Koffijberg, K. & C. Kowallik (2024): Ergebnisse der Gänsezählungen in Nordrhein-Westfalen im Juli 2024. NWO-Monitoringbericht 2024/01, Voerde/Duisburg. https://www.nw-ornithologen.de/images/textfiles/monitoring_downloads/Sommergaense_2024.pdf
- Koons, D. N., Rockwell, R. F., & Aubry, L. M. 2014. Effects of exploitation on an overabundant species: The lesser snow goose predicament. *Journal of Animal Ecology*, 83(2), 365–374.
- Lameris, T.K., O.B. Pokrovskaya, A.V. Kondratyev, Y.A. Anisimov, N.H. Buitendijk, P.M. Glazov, H.P. van der Jeugd, C. Kampichler, H. Kruckenberg, K.E. Litvin, J.A. Loshchagina, S. Moonen, G.J.D. Müskens, B.A. Nolet, K.H.T. Schreven, H. Sierdsema, E.M. Zaynagutdinova & M.P. Boom. 2023. Barnacle geese *Branta leucopsis* breeding on Novaya Zemlya: current distribution and population size estimated from tracking data. *Polar Biology* 46, 67–76. <https://doi.org/10.1007/s00300-022-03110-8>
- Madsen, J., G. Cracknell & A.D. Fox (editors). 1999. Goose Populations of the Western Palearctic: A Review of Status and Distribution. National Environmental Research Institute, Denmark and Wetlands International, Wageningen, The Netherlands. Wetlands International Publication Vol. 48.
- Madsen, J., Pedersen, C.L., Piironen, A., Liljebäck, N., Månsson, J. & Heinicke, T. (2025). Defining Favourable Reference Values for the Finland & NW Russia (br) Population of Taiga Bean Goose (*Anser fabalis fabalis*). EGMP Technical Report No. 28, Bonn, Germany. https://egmp.aewa.info/sites/default/files/download/population_status_reports/Draft_Report_Defining_FRVs_TB_G_Finland_NW_Russia.pdf
- Madsen, J., Sørensen, I.H., Johnson, F.A., Germain, R.R., Lewis, M., Leles, B.P. (Compilers) 2025. AEWA International Single Species Management Plan for the Pink-footed Goose (*Anser brachyrhynchus*) – Svalbard breeding population. AEWA Technical Series No. 79. Bonn, Germany.
- Marescot, L., G. Chapron, I. Chads, P. L. Fackler, C. Duchamp, E. Marboutin, and O. Gimenez. 2013. Complex decisions made simple: a primer on stochastic dynamic programming. *Methods in Ecology and Evolution*, 4: 872–884, <https://doi.org/10.1111/2041-210X.12082>
- McIntosh, A. L. S., S. Bearhop, G. M. Hilton, J. M. Shaw, and F. A. Johnson. 2023. Modelling harvest of Greenland barnacle geese and its implications in mitigating human-wildlife conflict. *Journal of Applied Ecology* 60:764–777.
- Månsson, J., N. Liljebäck, R. Buij, S. Moonen, and J. Elmberg (2024). Regional differences in crippling rate in greylag geese *Anser anser*. *Wildlife Biology* (early view). <https://doi.org/10.1002/wlb3.01356>.

Nagy, S., H. Heldbjerg, G. H. Jensen, F. A. Johnson, J. Madsen, E. Meyers, and S. Dereliev. 2021a. Adaptive Flyway Management Programme East Greenland/Scotland & Ireland Population of the Barnacle Goose (*Branta leucopsis*). AEWA EGMP Programme No. 2. Bonn, Germany.

Nagy, S., H. Heldbjerg, G. H. Jensen, F. A. Johnson, J. Madsen, O. Therkildsen, E. Meyers and S. Dereliev, 2021b. Adaptive Flyway Management Programme for Russia/Germany & Netherlands Population of the Barnacle Goose (*Branta leucopsis*). AEWA EGMP Programme No. 3. Bonn, Germany.

Nagy, S., H. Heldbjerg, G. H. Jensen, F. A. Johnson, and J. Madsen. 2021c. Adaptive Flyway Management Programme for the Greylag Goose (*Anser anser*), NW Europe/SW Europe population. AEWA EGMP Programme No. 1. Bonn, Germany.

Niedersächsische Sommer-Gänsezählung 2025.

https://www.nlwkn.niedersachsen.de/sommergaense/methode_und_bearbeitungsintestat/sommer-gansezahlungen-in-niedersachsen-methode-und-bearbeitungsintensitat-244297.html

Nielsen, R.D., Holm, T.E., Clausen, P., Bregnballe, T., Clausen, K.K., Petersen, I.K., Sterup, J., Balsby, T.J.S., Pedersen, C.L., Dalby, L., Mikkelsen, P., Møllerup, K.A. & Bladt J. 2023. Fugle 2020-2021. NOVANA. Aarhus Universitet, DCE – Nationalt Center for Miljø og Energi. Videnskabelig rapport fra DCE - Nationalt Center for Miljø og Energi nr. 531. <http://novana.au.dk/fugle/>

Nielsen, R.D., T.E. Holm, P. Clausen, J. Sterup, C.L. Pedersen, K.K. Clausen, T. Bregnballe, H.M. Thomsen, T.J.S. Balsby, I.K. Petersen, I.K., P. Mikkelsen, L. Dalby, and K.A. Møllerup. 2024. Fugle 2018-2023. NOVANA. Aarhus Universitet, DCE – Nationalt Center for Miljø og Energi. Videnskabelig rapport nr. 633. <https://novana.au.dk/fugle/2018-2023>.

Noer, H., J. Madsen, and P. Hartmann. 2007. Reducing wounding of game by shotgun hunting: Effects of a Danish action plan on pink-footed geese. *Journal of Applied Ecology*, 44: 653-62, <https://doi.org/10.1111/j.1365-2664.2007.01293.x>

Onkelinx, T., K. Devos, and P. Quataert. 2017. Working with population totals in the presence of missing data. Comparing imputation methods in terms of bias and precision. *Journal of Ornithology*, 158, 603-615, <https://doi.org/10.1007/s10336-016-1404-9>

Piironen, A., A.D. Fox, H. Kampe-Persson, U. Skjellberg, O.R. Therkildsen, and T. Laaksonen. 2023. ‘When and where to count? Implications of migratory connectivity and nonbreeding distribution to population censuses in a migratory bird population’. *Population Ecology* 65:121–132.

Pistorius, P.A., A. Follestad and F.E. Taylor. 2006. Declining winter survival and fitness implications associated with latitudinal distribution in Norwegian Greylag Geese *Anser anser*. *Ibis* 148(1):114-125, <https://doi.org/10.1111/j.1474-919X.2006.00498.x>

Schaub, M., and F. Abadi. 2011. Integrated population models: a novel analysis framework for deeper insights into population dynamics. *Journal of Ornithology*, 152(Suppl 1):227-37, <https://doi.org/10.1007/s10336-010-0632-7>

Schneider, E., and L. Bacon. 2022. Etude démographique des oies cendrées sur la voie de migration européenne NO/SO. 2e année de Master Ecologie-Ethologie, 2021-2022. Université Jean Monnet, Saint-Etienne.

Sedinger, J. S., Nicolai, C. A., Lensink, C. J. Wentworth, C., & B. Conant. 2007. Black Brant harvest, density dependence, and survival: A record of population dynamics. *Journal of Wildlife Management*, 71(2), 496–506.

Sørensen, I.H., F.A. Johnson, R.R. Germain & J. Madsen (2024): Estimating Greylag Goose breeding population size and productivity. Status and recommendations for post-breeding population counts and age ratio surveys in breeding range states of Management Unit 1 of the NW Europe (br) population of Greylag Goose: Denmark, Norway, Sweden, and Finland. EGMP Technical Report No. 23, Bonn, Germany.

Sørensen, I.H., R.R. Germain, F.A. Johnson, H. Baveco, K. Koffijberg, and J. Madsen. 2025. Population Status and Assessment Report 2025. EGMP Technical Report No. 26. Bonn, Germany

Sovon. 2022. Advies voor de uitvoering van het ganzenbeheer in de provincie Noord-Holland - onderdeel rekenkundige benadering ondergrens populatiebeheer. Sovon-notitie 2022/14. Sovon Vogelonderzoek Nederland, Nijmegen.

Sovon. 2025. Vogelbalans 2025. Sovon Vogelonderzoek Nederland, Nijmegen, <https://stats.sovon.nl/pub/publicatie/22635>

van Roomen, M., Citegetse, G., Crowe, O., Dodman, T., Hagemeyer, W., Meise, K., & Schekkerman, H. (eds.). 2025. East Atlantic Flyway Assessment 2023. The status of coastal waterbird populations and their sites. Wadden Sea Flyway Initiative p/a CWSS, Wilhelmshaven, Germany, Wetlands International, Wageningen, The Netherlands, BirdLife International, Cambridge, United Kingdom. <https://doi.org/10.5281/zenodo.15355686>

Volkov, S.V. and S. Rozenfeld. 2025. Remote Tracking of the Spatio-Temporal Distribution of Western Taiga Bean Geese (*Anser fabalis fabalis*, Anseriformes, Aves) in Russia for Improving the Conservation Strategy. *Biology Bulletin* Vol. 52:411.

Wolff S., B. Koop, H.-J. Augst, K.Günther and Uwe Helbing. 2023. Rastbestände von Weißwangen- und Graugänsen 2022. In: Jahresbericht 2023 Zur biologischen Vielfalt. Jagd und Artenschutz, Schleswig-Holstein. https://www.schleswig-holstein.de/DE/fachinhalte/A/artenschutz/Downloads/jahresbericht2023.pdf?__blob=publicationFile&v=6.

Wylegała, P., and B. Smyk. 2025. Goose Monitoring. In: Chodkiewicz T., Przymencki M. (eds.) 2025. Report on fieldwork and the compilation of results obtained during the migration and wintering period in the 2024/2025 season. Bird monitoring including Natura 2000 Special Protection Areas, years 2023-2025. Stage 5. Task 1. Bird monitoring - fieldwork and results compilation. GIOŚ, Warsaw.