

Range State: Norway



Ingunn Tombre

Where do the damage occur?

When does it occur?

What is damaged?

Who is doing the damage (culprit species)?



Who is doing the damage

Pink-footed Goose




Barnacle Goose




Greylag Goose



Where do the damage occur?



Svalbard-breeding
Pink-footed Goose & **Barnacle Goose**
(Arctic)




Where do the damage occur?

Greylag Geese

Breeding observations
from the online
registration platform
www.artsobservasjoner.no

At present –
the challenges are
on farmland...



Where do the damage occur?

Temperate-breeding **Barnacle** Geese



The challenges are on farmland...



When does it occur?

Norway is a country with a long winter...

The main agriculture challenges with geese are in **SPRING**

But... also during the rest of the growing season (summer, early autumn)



Christiane Hübner

When does it occur?

Mid- and North-Norway

Early spring:

The growing grass is fragile and important for the farmers, especially for the **sheep farmers** who need the grass for the new-born lambs



When does it occur?

Mid-Norway

Early spring;
New-sown cereals



When does it occur?

Middle and southern parts of Norway



Summer and **early autumn** situation challenging for cereal and vegetable producers




What is damaged?

Generally reflects the agriculture production and **distribution** in Norway



Cereals, pasture, vegetables
(lettuce, peas)

High costs!
Zero tolerance...




Summary for Norway

SPRING

Mid-Norway (Trøndelag)
North Norway (Helgeland & Vesterålen)

SPRING, SUMMER, AUTUMN

Oslofjord-area



«All over Norway» (coast)

SPRING, SUMMER, AUTUMN

- Subsidy scheme for the arctic-breeding geese
- No subsidies or compensations for greylags or Oslofjorden-barnies
- Tools in the tool-box!